

XII CBSE

**RELATION
FUNCTIONS
MATHEMATICS**



YOUR GATEWAY TO EXCELLENCE IN

IIT-JEE, NEET AND CBSE EXAMS

**RELATION
FUNCTIONS**

CONTACT US:



+91-9939586130
+91-9955930311



www.aepstudycircle.com



aepstudycircle@gmail.com



2ND FLOOR, SATKOURI COMPLEX, THANA CHOWK, RAMGARH - 829122-JH

RELATION FUNCTION

BASIC CONCEPTS

- 1. Relation:** If A and B are two non-empty sets, then any subset R of $A \times B$ is called relation from set A to set B .

i.e., $R : A \rightarrow B \Leftrightarrow R \subseteq A \times B$

If $(x, y) \in R$, then we write $x R y$ (read as x is R related to y) and if $(x, y) \notin R$, then we write $x \not R y$ (read as x is not R related to y).

- 2. Domain and Range of a Relation:** If R is any relation from set A to set B then,

(a) **Domain of R** is the set of all first coordinates of elements of R and it is denoted by $\text{Dom}(R)$.

(b) **Range of R** is the set of all second coordinates of R and it is denoted by $\text{Range}(R)$.

A relation R on set A means, the relation from A to A i.e., $R \subseteq A \times A$.

- 3. Some Standard Types of Relations:**

Let A be a non-empty set. Then, a relation R on set A is said to be

(a) **Reflexive:** If $(x, x) \in R$ for each element $x \in A$, i.e., if xRx for each element $x \in A$.

(b) **Symmetric:** If $(x, y) \in R \Rightarrow (y, x) \in R$ for all $x, y \in A$, i.e., if $xRy \Rightarrow yRx$ for all $x, y \in A$.

(c) **Transitive:** If $(x, y) \in R$ and $(y, z) \in R \Rightarrow (x, z) \in R$ for all $x, y, z \in A$, i.e., if xRy and $yRz \Rightarrow xRz$.

- 4. Equivalence Relation:** Any relation R on a set A is said to be an equivalence relation if R is reflexive, symmetric and transitive.

- 5. Antisymmetric Relation:** A relation R in a set A is antisymmetric

if $(a, b) \in R, (b, a) \in R \Rightarrow a = b \forall a, b \in R$, or aRb and $bRa \Rightarrow a = b, \forall a, b \in R$.

For example, the relation "greater than or equal to, \geq " is antisymmetric relation as

$$a \geq b, b \geq a \Rightarrow a = b \forall a, b$$

[Note: "Antisymmetric" is completely different from not symmetric.]

- 6. Equivalence Class:** Let R be an equivalence relation on a non-empty set A . For all $a \in A$, the equivalence class of ' a ' is defined as the set of all such elements of A which are related to ' a ' under R . It is denoted by $[a]$.

i.e., $[a] = \text{equivalence class of } 'a' = \{x \in A : (x, a) \in R\}$

- 7. Function:** Let X and Y be two non-empty sets. Then, a rule f which associates to each element $x \in X$, a unique element, denoted by $f(x)$ of Y , is called a function from X to Y and written as $f : X \rightarrow Y$ where, $f(x)$ is called image of x and x is called the **pre-image** of $f(x)$ and the set Y is called the **co-domain** of f and $f(X) = \{f(x) : x \in X\}$ is called the range of f .

8. Types of Function:

(i) **One-one function (injective function):** A function $f: X \rightarrow Y$ is defined to be one-one if the image of distinct element of X under rule f are distinct, i.e., for every $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies that $x_1 = x_2$.


(ii) **Onto function (Surjective function):** A function $f: X \rightarrow Y$ is said to be onto function if each element of Y is the image of some element of X i.e., for every $y \in Y$, there exists some $x \in X$, such that $y = f(x)$. Thus f is onto if range of $f =$ co-domain of f .

(iii) **One-one onto function (Bijective function):** A function $f: X \rightarrow Y$ is said to be one-one onto, if f is both one-one and onto.

(iv) **Many-one function:** A function $f: X \rightarrow Y$ is said to be a many-one function if two or more elements of set X have the same image in Y . i.e.,

$f: X \rightarrow Y$ is a many-one function if there exist $a, b \in X$ such that $a \neq b$ but $f(a) = f(b)$.

9. **Composition of Functions:** Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be two functions. Then, the composition of f and g , denoted by $g \circ f$, is defined as the function.



$$g \circ f: A \rightarrow C \quad \text{given by}$$

$$g \circ f(x) = g(f(x)), \forall x \in A$$

Clearly, $\text{dom}(g \circ f) = \text{dom}(f)$

Also, $g \circ f$ is defined only when $\text{range}(f) \subseteq \text{dom}(g)$

10. **Identity Function:** Let R be the set of real numbers. A function $I: R \rightarrow R$ such that

$$I(x) = x \quad \forall x \in R \text{ is called identity function.}$$

Obviously, identity function associates each real number to itself.

11. **Invertible Function:** For $f: A \rightarrow B$, if there exists a function $g: B \rightarrow A$ such that $g \circ f = I_A$ and $f \circ g = I_B$, where I_A and I_B are identity functions, then f is called an invertible function, and g is called the inverse of f and it is written as $f^{-1} = g$.

12. **Number of Functions:** If X and Y are two finite sets having m and n elements respectively then the number of functions from X to Y is n^m .

13. **Vertical Line Test:** It is used to check whether a relation is a function or not. Under this test, graph of given relation is drawn assuming elements of domain along x -axis. If a vertical line drawn anywhere in the graph, intersects the graph at only one point then the relation is a function, otherwise it is not a function.

14. **Horizontal Line Test:** It is used to check whether a function is one-one or not. Under this test graph of given function is drawn assuming elements of domain along x -axis. If a horizontal line (parallel to x -axis) drawn anywhere in graph, intersects the graph at only one point then the function is one-one, otherwise it is many-one.

MULTIPLE CHOICE QUESTIONS

Choose and write the correct option in the following questions.

- The relation R in the set $A = \{1, 2, 3, 4\}$ given by $R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$ is
 (a) reflexive and symmetric but not transitive
 (b) reflexive and transitive but not symmetric
 (c) symmetric and transitive but not reflexive
 (d) an equivalence relation
- If $A = \{a, b, c, d\}$, then a relation $R = \{(a, b), (b, a), (a, a)\}$ on A is
 (a) symmetric only
 (b) transitive only
 (c) reflexive and transitive
 (d) symmetric and transitive only
- For real numbers x and y , define xRy if and only if $x - y + \sqrt{2}$ is an irrational number. Then the relation R is [NCERT Exemplar]
 (a) reflexive
 (b) symmetric
 (c) transitive
 (d) none of these
- Consider the non-empty set consisting of children in a family and a relation R defined as aRb if a is brother of b . Then R is [NCERT Exemplar]
 (a) symmetric but not transitive
 (b) transitive but not symmetric
 (c) neither symmetric nor transitive
 (d) both symmetric and transitive
- The maximum number of equivalence relation on the set $A = \{1, 2, 3\}$ are [NCERT Exemplar]
 (a) 1
 (b) 2
 (c) 3
 (d) 5
- Let L denotes the set of all straight lines in a plane. Let a relation R be defined by lRm if and only if l is perpendicular to $m \forall l, m \in L$. Then R is [NCERT Exemplar]
 (a) reflexive
 (b) symmetric
 (c) transitive
 (d) none of these
- Let $A = \{1, 2, 3\}$. Then number of relations containing $(1, 2)$ and $(1, 3)$ which are reflexive and symmetric but not transitive is
 (a) 1
 (b) 2
 (c) 3
 (d) 4
- Let $A = \{1, 2, 3\}$. Then number of equivalence relations containing $(1, 2)$ is/are
 (a) 1
 (b) 2
 (c) 3
 (d) 4
- Let A and B be finite sets containing m and n elements respectively. The number of relations that can be defined from A to B is
 (a) 2^{mn}
 (b) 2^{m+n}
 (c) mn
 (d) 0
- Set A has 3 elements and the set B has 4 elements. Then the number of injective mapping that can be defined from A to B is [NCERT Exemplar]
 (a) 144
 (b) 12
 (c) 24
 (d) 64
- The function $f: R \rightarrow R$ defined by $f(x) = 2^x + 2^{|x|}$ is
 (a) One-one and onto
 (b) Many-one and onto
 (c) One-one and into
 (d) Many-one and into
- If the set A contains 5 elements and the set B contains 6 elements, then the number of one-one and onto mapping from A to B is
 (a) 720
 (b) 120
 (c) 0
 (d) none of these
- Which of the following functions from Z into Z is bijection? [NCERT Exemplar]
 (a) $f(x) = x^3$
 (b) $f(x) = x + 2$
 (c) $f(x) = 2x + 1$
 (d) $f(x) = x^2 + 1$
- Let $f: [2, \infty) \rightarrow R$ be the function defined by $f(x) = x^2 - 4x + 5$, then the range of f is [NCERT Exemplar]
 (a) R
 (b) $[1, \infty)$
 (c) $[4, \infty)$
 (d) $[5, \infty)$

15. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = x^2 + 1$. Then, pre-images of 17 and -3, respectively, are [NCERT Exemplar]
 (a) $\phi, \{4, -4\}$ (b) $\{3, -3\}, \phi$ (c) $\{4, -4\}, \phi$ (d) $\{4, -4\}, \{2, -2\}$
16. Let the function $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = 2x + \sin x$ for $x \in \mathbb{R}$. Then f is
 (a) one-one but not onto (b) onto but not one-one
 (c) neither one-one nor onto (d) one-one and onto
17. Let R be the relation in the set \mathbb{N} given by $R = \{(a, b) : a = b - 2, b > 6\}$ choose the correct answer.
 (a) $(2, 4) \in R$ (b) $(3, 8) \in R$ (c) $(6, 8) \in R$ (d) $(8, 7) \in R$
18. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x^4$. Choose the correct answer
 (a) f is one-one onto (b) f is many one onto
 (c) f is one-one but not onto (d) f is neither one-one nor onto.
19. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 3x$. Choose the correct answer.
 (a) f is one-one onto. (b) f is many one onto.
 (c) f is one-one but not onto (d) f is neither one-one nor onto.
20. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 2x^3 + 2x^2 + 300x + 5 \sin x$ then f is
 (a) one-one onto (b) one-one into (c) many one onto (d) many one into
21. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = x^2 + 1$. Then, pre-image of 5 and -5, respectively are
 (a) $\phi, \{-2\}$ (b) $\{(3, -3)\}, \phi$ (c) $\{-2, 2\}, \phi$ (d) $\{1, -1\}, \{2, -2\}$
22. The domain of the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \sqrt{x^2 - 4}$ is
 (a) $[-2, 2]$ (b) $(-2, 2)$ (c) $(-\infty, -2] \cup [2, \infty)$ (d) $(-\infty, \infty)$
23. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} 3x, & \text{if } x > 3 \\ x^2, & \text{if } 1 < x \leq 3 \\ x, & \text{if } x \leq 1 \end{cases}$
 Then $f(-2) + f(0) + f(2) + f(5)$ is equal to
 (a) 0 (b) 17 (c) -4 (d) none of these
24. Let R is reflexive relation on a finite set A having n element, and let there be m ordered pairs in R . Then
 (a) $m \geq n$ (b) $m \leq n$ (c) $m = n$ (d) none of these
25. The domain of the function $f(x) = \log_{3+x}(x^2 - 1)$ is
 (a) $(-3, -1) \cup (1, \infty)$ (b) $[-3, -1] \cup [1, \infty)$
 (c) $(-3, -2) \cup (-2, -1) \cup (1, \infty)$ (d) $[-3, -2) \cup (-2, -1) \cup [1, \infty)$
26. Let $f: \mathbb{R} \rightarrow [0, \frac{\pi}{2})$ defined by $f(x) = \tan^{-1}(x^2 + x + a)$, then the set of values of a for which f is onto is
 (a) $[0, \infty)$ (b) $[\frac{1}{4}, \infty)$ (c) $[2, 1]$ (d) none of these
27. If the function $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ are defined as

$$f(x) = \begin{cases} 0, & x \in \text{rational} \\ x, & x \in \text{irrational} \end{cases}$$
 and
$$g(x) = \begin{cases} 0, & x \in \text{irrational} \\ x, & x \in \text{rational} \end{cases}$$
 then $(f - g)$ is
 (a) one-one onto. (b) many-one onto
 (c) one-one but not onto (d) neither one-one nor onto.

28. If a relation R on the set $\{1, 2, 3, 4\}$ is defined by $R = \{(1, 2), (3, 4)\}$. Then R is
(a) reflexive (b) transitive (c) symmetric (d) none of these
29. If the set A contains 4 elements and the set B contains 5 elements, then the number of one-one and onto mappings from A to B is
(a) 0 (b) 4^5 (c) 5^4 (d) none of these
30. Let $A = \{x, y, z\}$ and $B = \{a, b\}$ then the number of onto function from A to B is
(a) 0 (b) 3 (c) 6 (d) 8
31. If A and B have 4 and 6 elements respectively then the number of one-one function from A to B is
(a) 4^6 (b) 6^4 (c) 360 (d) 240
32. If A and B have 4 elements each then the number of one-one onto (bijective) function from A to B is
(a) 0 (b) 24 (c) 4^2 (d) None of these
33. If R is an equivalence relation on A , then R^{-1} on A is
(a) Transitive only (b) Symmetric only (c) Reflexive only (d) Equivalence relation
34. The relation "greater than" denoted by $>$ in the set of integers is
(a) Symmetric (b) Reflexive (c) Transitive (d) None of these
35. If R_1 and R_2 are symmetric relations in a set A , then $R_1 \cup R_2$ is
(a) Reflexive (b) Symmetric (c) Transitive (d) None of these
36. The function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 4^x + 4^{|x|}$ is
(a) one-one and into (b) one-one and onto
(c) many one and into (d) many one and onto
37. Identity relation R on a set A is
(a) Reflexive only (b) Symmetric only (c) Transitive only (d) Equivalence
38. The relation "congruence modulo m " on the set \mathbb{Z} of all integers is a relation of type
(a) Reflexive only (b) Symmetric only (c) Transitive only (d) Equivalence
39. Let $f: \mathbb{R} \rightarrow \left[0, \frac{\pi}{2}\right)$ defined by $f(x) = \tan^{-1}(x^2 + x + 2a)$ then the set of values of ' a ' for which f is onto, is
(a) $\left(-\frac{1}{4}, \infty\right)$ (b) $[-1, \infty)$ (c) $\left[-\frac{1}{8}, \infty\right)$ (d) $\left[\frac{1}{8}, \infty\right)$
40. If the function $f(x)$ satisfying $(f(x))^2 - 4f(x)f'(x) + (f'(x))^2 = 0$ then $f(x)$ equals
(a) $\lambda e^{(2+\sqrt{5})x}$ (b) $\lambda e^{(2-\sqrt{5})x}$ (c) $\lambda e^{(2\pm\sqrt{3})x}$ (d) $\lambda e^{(3-\sqrt{3})x}$
41. Let $f: (-1, 1) \rightarrow B$ where $f(x) = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ is one-one and onto, then B equals
(a) $\left[0, \frac{\pi}{2}\right]$ (b) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (c) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (d) $\left(0, \frac{\pi}{2}\right)$
42. The function $y = \frac{x}{1+|x|}$, $x \in \mathbb{R}$, $y \in \mathbb{R}$ is
(a) One-one onto (b) Onto but not one-one
(c) One-one but not onto (d) None of these
43. A relation R in the set of non-zero complex number is defined by $z_1 R z_2 \Leftrightarrow \frac{z_1 - z_2}{z_1 + z_2}$ is real, then R is
(a) Reflexive (b) Symmetric (c) Transitive (d) Equivalence
44. Number of onto (subjective) functions from A to B if $n(A) = 6$ and $n(B) = 3$ are
(a) $2^6 - 2$ (b) $3^6 - 3$ (c) 340 (d) None of these

45. Let $A = \{7, 8, 9, 10\}$ and $R = \{(8, 8), (9, 9), (10, 10), (7, 8)\}$ be a relation on A , then R is
(a) Transitive (b) Reflexive (c) Symmetric (d) None of these
46. Let f, g be a function from the set $\{1, 2, \dots, 12\}$ to the set $\{1, 2, 3, \dots, 11\}$ then which of the following is correct?
(a) Number of onto functions from A to $B = \frac{12 \times 11}{2}$
(b) Total number of functions from A to $B = 11^{12}$
(c) The functions which are not onto $= 11^{12} - \frac{12 \times 11}{2}$
(d) All of these
47. Let p and q are positive integers, f is a function defined for positive numbers and attains only positive values such that $f[x f(y)] = x^a y^b$ then
(a) $b^2 = a$ (b) $a = b$ (c) $a^2 = b$ (d) None of these
48. Let $f: (-1, 1) \rightarrow B$, be a function defined by $f(x) = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ then f is both one-one and onto when B is the interval
(a) $\left[0, \frac{\pi}{2}\right)$ (b) $\left(0, \frac{\pi}{2}\right)$ (c) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
49. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \frac{e^{x^2} - e^{-x^2}}{e^{x^2} + e^{-x^2}}$ then
(a) $f(x)$ is one-one but not onto (b) $f(x)$ is neither one-one nor onto
(c) $f(x)$ is many one but onto (d) $f(x)$ is one-one and onto
50. If $A = \{7, 8, 9\}$, then the relation $R = \{(8, 9)\}$ in A is
(a) Symmetric only (b) Non-symmetric (c) Reflexive only (d) Equivalence
51. Let A be the finite set containing n distinct elements. The number of relations that can be defined on A is
(a) 2^n (b) n^2 (c) 2^{n^2} (d) 2^{n-1}
52. Let R_1 and R_2 be equivalence relations on a set A , then $R_1 \cup R_2$ may or may not be
(a) Reflexive (b) Symmetric (c) Transitive (d) None of these
53. Let R be the relation defined on the set N of natural numbers by the rule $x R y$ iff $x + 2y = 8$, then domain of R is
(a) $\{2, 4, 8\}$ (b) $\{2, 4, 6\}$ (c) $\{2, 4, 6, 8\}$ (d) $\{1, 2, 3, 4\}$
54. Let $A = \{a, b, c\}$ and $R = \{(a, a), (b, b), (c, c), (b, c), (a, b)\}$ be a relation on A , then R is
(a) Symmetric (b) Transitive (c) Reflexive (d) Equivalence
55. "Every relation is a function and every function is a relation" then which is correct for given statement?
(a) True (b) False (c) Can't say anything (d) None of these
56. If a relation R on the set $\{1, 2, 3\}$ be defined by $R = \{(1, 2)\}$, then R is
(a) Reflexive (b) Transitive (c) Symmetric (d) None of these
57. Let us define a relation R in R as $a R b$ if $a \geq b$. Then, R is
 $R = \{(a, b) : a \geq b\}$
(a) An equivalence relation
(b) Reflexive, transitive but not symmetric
(c) Symmetric, transitive but not reflexive
(d) Neither transitive nor reflexive but symmetric

58. If $A = \{1, 2, 3\}$ and consider the relation $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$. Then, R is
 (a) Reflexive but not symmetric (b) Reflexive but not transitive
 (c) Symmetric and transitive (d) Neither Symmetric nor transitive
59. The relation R defined on the set $A = \{1, 2, 3, 4, 5\}$ by $R = \{(a, b) : |a^2 - b^2| < 7\}$ is given by
 (a) $\{(1, 1), (2, 1), (3, 1), (4, 1), (2, 3)\}$
 (b) $\{(2, 2), (3, 2), (4, 2), (2, 4)\}$
 (c) $\{(3, 3), (4, 3), (5, 4), (3, 4)\}$
 (d) $\{(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (1, 2), (2, 3)\}$

Answers

- | | | | | | |
|---------|---------|---------|---------|---------|---------|
| 1. (b) | 2. (d) | 3. (a) | 4. (b) | 5. (d) | 6. (b) |
| 7. (a) | 8. (b) | 9. (a) | 10. (c) | 11. (c) | 12. (c) |
| 13. (b) | 14. (b) | 15. (c) | 16. (d) | 17. (c) | 18. (d) |
| 19. (a) | 20. (a) | 21. (c) | 22. (c) | 23. (b) | 24. (a) |
| 25. (c) | 26. (b) | 27. (a) | 28. (b) | 29. (a) | 30. (c) |
| 31. (c) | 32. (b) | 33. (d) | 34. (c) | 35. (b) | 36. (a) |
| 37. (d) | 38. (d) | 39. (d) | 40. (c) | 41. (c) | 42. (c) |
| 43. (d) | 44. (d) | 45. (a) | 46. (d) | 47. (c) | 48. (c) |
| 49. (b) | 50. (b) | 51. (c) | 52. (c) | 53. (b) | 54. (c) |
| 55. (b) | 56. (b) | 57. (b) | 58. (a) | 59. (d) | |

CASE-BASED QUESTIONS

Choose and write the correct option in the following questions.

1. Read the following and answer any four questions from (i) to (v).

A general election of Lok Sabha is a gigantic exercise. About 911 million people were eligible to vote and voter turnout was about 67%, the highest ever



Let I be the set of all citizens of India who were eligible to exercise their voting right in general election held in 2019. A relation ' R ' is defined on I as follows:

$$R = \{(V_1, V_2) : V_1, V_2 \in I \text{ and both use their voting right in general election - 2019}\}$$

[CBSE Question Bank]

Answer the questions given below.

- (i) Two neighbours X and $Y \in I$. X exercised his voting right while Y did not cast her vote in general election – 2019. Which of the following is true?
- (a) $(X, Y) \in R$ (b) $(Y, X) \in R$
(c) $(X, X) \notin R$ (d) $(X, Y) \notin R$
- (ii) Mr. 'X' and his wife 'W' both exercised their voting right in general election -2019, Which of the following is true?
- (a) both (X, W) and $(W, X) \in R$ (b) $(X, W) \in R$ but $(W, X) \notin R$
(c) both (X, W) and $(W, X) \notin R$ (d) $(W, X) \in R$ but $(X, W) \notin R$
- (iii) Three friends F_1, F_2 and F_3 exercised their voting right in general election- 2019, then which of the following is true?
- (a) $(F_1, F_2) \in R, (F_2, F_3) \in R$ and $(F_1, F_3) \in R$
(b) $(F_1, F_2) \in R, (F_2, F_3) \in R$ and $(F_1, F_3) \notin R$
(c) $(F_1, F_2) \in R, (F_2, F_3) \in R$ but $(F_3, F_3) \notin R$
(d) $(F_1, F_2) \notin R, (F_2, F_3) \notin R$ and $(F_1, F_3) \notin R$
- (iv) The above defined relation R is
- (a) Symmetric and transitive but not reflexive
(b) Universal relation
(c) Equivalence relation
(d) Reflexive but not symmetric and transitive
- (v) Mr. Shyam exercised his voting right in General Election – 2019, then Mr. Shyam is related to which of the following?
- (a) All those eligible voters who cast their votes
(b) Family members of Mr. Shyam
(c) All citizens of India
(d) Eligible voters of India

Sol. We have a relation ' R ' is defined on I as follows:

$$R = \{V_1, V_2\} : V_1, V_2 \in I \text{ and both use their voting right in general election – 2019}$$

- (i) Two neighbors X and $Y \in I$. Since X exercised his voting right while Y did not cast her vote in general election – 2019
Therefore, $(X, Y) \notin R$
 \therefore Option (d) is correct.
- (ii) Since Mr. 'X' and his wife 'W' both exercised their voting right in general election – 2019.
 \therefore Both (X, W) and $(W, X) \in R$.
 \therefore Option (a) is correct.
- (iii) Since three friends F_1, F_2 and F_3 exercised their voting right in general election – 2019, therefore
 $(F_1, F_2) \in R, (F_2, F_3) \in R$ and $(F_1, F_3) \in R$
 \therefore Option (a) is correct.
- (iv) This relation is an equivalence relation
 \therefore Option (c) is correct.
- (v) Mr. Shyam exercised his voting right in General election – 2019, then Mr. Shyam is related to all those eligible votes who cast their votes.
 \therefore Option (a) is correct.

2. Read the following and answer any four questions from (i) to (v).

Sherlin and Danju are playing Ludo at home during Covid-19. While rolling the dice, Sherlin's sister Raji observed and noted the possible outcomes of the throw every time belongs to set $\{1,2,3,4,5,6\}$. Let A be the set of players while B be the set of all possible outcomes.



$$A = \{S, D\}, B = \{1,2,3,4,5,6\}$$

[CBSE Question Bank]

Answer the questions given below.

(i) Let $R : B \rightarrow B$ be defined by $R = \{(x, y) : y \text{ is divisible by } x\}$ is

- (a) Reflexive and transitive but not symmetric
- (b) Reflexive and symmetric and not transitive
- (c) Not reflexive but symmetric and transitive
- (d) Equivalence

(ii) Raji wants to know the number of functions from A to B . How many number of functions are possible?

- (a) 6^2
- (b) 2^6
- (c) $6!$
- (d) 2^{12}

(iii) Let R be a relation on B defined by $R = \{(1,2), (2,2), (1,3), (3,4), (3,1), (4,3), (5,5)\}$. Then R is

- (a) Symmetric
- (b) Reflexive
- (c) Transitive
- (d) None of these three

(iv) Raji wants to know the number of relations possible from A to B . How many numbers of relations are possible?

- (a) 6^2
- (b) 2^6
- (c) $6!$
- (d) 2^{12}

(v) Let $R : B \rightarrow B$ be defined by $R = \{(1,1), (1,2), (2,2), (3,3), (4,4), (5,5), (6,6)\}$, then R is

- (a) Symmetric
- (b) Reflexive and Transitive
- (c) Transitive and symmetric
- (d) Equivalence

Sol. (i) Given $R : B \rightarrow B$ be defined by

$$R = \{(x, y) : y \text{ is divisible by } x\}$$

Reflexive : Let $x \in B$, since x always divide x itself.

$$\therefore (x, x) \in R$$

It is reflexive.

Symmetric : Let $x, y \in B$ and let $(x, y) \in R$

$$\Rightarrow y \text{ is divisible by } x$$

$$\Rightarrow \frac{y}{x} = k_1, \text{ where } k_1 \text{ is an integer.}$$

$$\Rightarrow \frac{x}{y} = \frac{1}{k_1} \neq \text{integer.}$$

$$\therefore (y, x) \notin R$$

It is not symmetric.

Transitive : Let $x, y, z \in B$ and

$$\text{Let } (x, y) \in R \Rightarrow \frac{y}{x} = k_1, \text{ where } k_1 \text{ is an integer.}$$

$$\text{and, } (y, z) \in R \Rightarrow \frac{z}{y} = k_2, \text{ where } k_2 \text{ is an integer.}$$

$$\therefore \frac{y}{x} \times \frac{z}{y} = k_1 \cdot k_2 = k \text{ (integer)}$$

$$\Rightarrow \frac{z}{x} = k \Rightarrow (x, z) \in R$$

It is transitive.

Hence, relation is reflexive and transitive but not symmetric.

\therefore Option (a) is correct.

(ii) We have,

$$A = \{S, D\} \Rightarrow n(A) = 2$$

$$\text{and, } B = \{1, 2, 3, 4, 5, 6\} \Rightarrow n(B) = 6$$

\therefore Number of functions from A to B is 6^2 .

\therefore Option (a) is correct.

(iii) Given,

R be a relation on B defined by

$$R = \{(1, 2), (2, 2), (1, 3), (3, 4), (3, 1), (4, 3), (5, 5)\}$$

R is not reflexive since $(1, 1), (3, 3), (4, 4) \notin R$

R is not symmetric as $(1, 2) \in R$ but $(2, 1) \notin R$

and, R is not transitive as $(1, 3) \in R$ and $(3, 1) \in R$ but $(1, 1) \notin R$

$\therefore R$ is neither reflexive nor symmetric nor transitive.

\therefore Option (d) is correct.

(iv) Total number of possible relations from A to $B = 2^{12}$

\therefore Option (d) is correct.

(v) Given $R : B \rightarrow B$ be defined by $R = \{(1, 1), (1, 2), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}$

$\therefore R$ is reflexive as each elements of B is related to itself and R is also transitive as $(1, 2) \in R$ and $(2, 2) \in R$

$$\Rightarrow (1, 2) \in R$$

$\therefore R$ is reflexive and transitive.

\therefore Option (b) is correct.

3. Read the following and answer any four questions from (i) to (v).

An organization conducted bike race under 2 different categories-boys and girls. In all, there were 250 participants. Among all of them finally three from Category 1 and two from Category 2 were selected for the final race. Ravi forms two sets B and G with these participants for his college project.

Let $B = \{b_1, b_2, b_3\}$ $G = \{g_1, g_2\}$ where B represents the set of boys selected and G the set of girls who were selected for the final race.

[CBSE Question Bank]



Ravi decides to explore these sets for various types of relations and functions

Answer the questions given below.

- (i) Ravi wishes to form all the relations possible from B to G . How many such relations are possible?
 (a) 2^6 (b) 2^5 (c) 0 (d) 2^3
- (ii) Let $R : B \rightarrow B$ be defined by $R = \{(x, y) : x \text{ and } y \text{ are students of same sex}\}$, Then this relation R is
 (a) Equivalence
 (b) Reflexive only
 (c) Reflexive and symmetric but not transitive
 (d) Reflexive and transitive but not symmetric
- (iii) Ravi wants to know among those relations, how many functions can be formed from B to G ?
 (a) 2^2 (b) 2^{12} (c) 3^2 (d) 2^3
- (iv) Let $R : B \rightarrow G$ be defined by $R = \{(b_1, g_1), (b_2, g_2), (b_3, g_1)\}$, then R is
 (a) Injective (b) Surjective
 (c) Neither Surjective nor Injective (d) Surjective and Injective
- (v) Ravi wants to find the number of injective functions from B to G . How many numbers of injective functions are possible?
 (a) 0 (b) $2!$ (c) $3!$ (d) $0!$

Sol. We have sets

$$B = \{b_1, b_2, b_3\}, G = \{g_1, g_2\}$$

$$\Rightarrow n(B) = 3 \text{ and } n(G) = 2$$

(i) Number of all possible relations from B to $G = 2^{3 \times 2} = 2^6$

\therefore Option (a) is correct.

(ii) Given relation $R = \{(x, y) : x \text{ and } y \text{ are student of same sex}\}$

On the set B .

Since the set is $B = \{b_1, b_2, b_3\}$ = all boys

\therefore It is an equivalence relation.

\therefore Option (a) is correct.

(iii) We have,

$$B = \{b_1, b_2, b_3\} \Rightarrow n(B) = 3$$

$$G = \{g_1, g_2\} \Rightarrow n(G) = 2$$

\therefore Total no. of possible functions from B to $G = 2^3$

\therefore Option (d) is correct.

(iv) We have,

$R : B \rightarrow G$ be defined by

$$R = \{(b_1, g_1), (b_2, g_2), (b_3, g_1)\}$$

It is not injective because $(b_1, g_1) \in R$ and $(b_3, g_1) \in R$

So $b_1 \neq b_3 \Rightarrow$ same image g_1 .

It is surjective because its Co-domain = Range.

$\therefore R$ is Surjective.

\therefore Option (b) is correct.

(v) Since R is not injective therefore number of injective functions = 0

\therefore Option (a) is correct.

ASSERTION-REASON QUESTIONS

In the following questions, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
- (b) Both A and R are true but R is not the correct explanation of A.
- (c) A is true but R is false.
- (d) A is false and R is also false.

1. **Assertion (A):** Let L be the collection of all lines in a plane and R_1 be the relation on L as $R_1 = \{(L_1, L_2) : L_1 \perp L_2\}$ is a symmetric relation.

Reason (R): A relation R is said to be symmetric if $(a, b) \in R \Rightarrow (b, a) \in R$.

2. **Assertion (A):** Let R be the relation on the set of integers Z given by $R = \{(a, b) : 2 \text{ divides } (a - b)\}$ is an equivalence relation.

Reason (R): A relation R in a set A is said to be an equivalence relation if R is reflexive, symmetric and transitive.

3. **Assertion (A):** Let $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x$, then f is a one-one function.

Reason (R): A function $g : A \rightarrow B$ is said to be onto function if for each $b \in B, \exists a \in A$ such that $g(a) = b$.

4. **Assertion (A):** Let function $f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ be an onto function. Then it must be one-one function.

Reason (R): A one-one function $g : A \rightarrow B$, where A and B are finite set and having same number of elements, then it must be onto and vice-versa.

Answers

1. (a) 2. (a) 3. (b) 4. (a)

HINTS/SOLUTIONS OF SELECTED MCQS

1. Since every element of A is related to itself in the given relation R , therefore R is reflexive and as $(1, 2) \in R$ and $(2, 2) \in R \Rightarrow (1, 2) \in R$ also $(1, 3) \in R$ and $(3, 2) \in R \Rightarrow (1, 2) \in R$. Again $(1, 3) \in R$ and $(3, 3) \in R \Rightarrow (1, 3) \in R$. Thus R is also transitive. Hence relation R is reflexive and transitive but not symmetric because, $(1, 2) \in R$ but $(2, 1) \notin R$, also $(1, 3) \in R$ but $(3, 1) \notin R$ and $(3, 2) \in R$ but $(2, 3) \notin R$.

Option (b) is correct.

2. On the set $A = \{a, b, c, d\}$ given relation $R = \{(a, b), (b, a), (a, a)\}$ is symmetric and transitive only.

Since, $(a, b) \in R \Rightarrow (b, a) \in R$, therefore it is symmetric

Also, $(a, b) \in R$ and $(b, a) \in R \Rightarrow (a, a) \in R$, so it is also transitive. As (b, b) , (c, c) and (d, d) does not belong to R hence R is not reflexive.

Hence relation R is symmetric and transitive only.

Option (d) is correct.

3. For any $x \in \mathbb{R}$

$$x - x + \sqrt{2} = \sqrt{2} \text{ is an irrational number} \Rightarrow (x, x) \in R \quad \forall x \in \mathbb{R}$$

$\therefore R$ is reflexive

For 2, $\sqrt{2} \in \mathbb{R}$

$$\sqrt{2} - 2 + \sqrt{2} = 2\sqrt{2} - 2 \text{ is an irrational number.}$$

$$\Rightarrow (\sqrt{2}, 2) \in R$$

$$\text{But } 2 - \sqrt{2} + \sqrt{2} = 2 \text{ which is a rational number}$$

$$\Rightarrow (2, \sqrt{2}) \notin R$$

$\Rightarrow R$ is not reflexive

R is not transitive

For 2, $\sqrt{3}, \sqrt{2} \in \mathbb{R}$

$$\because 2 - \sqrt{3} + \sqrt{2} = 2 - (\sqrt{3} - \sqrt{2}) \text{ is an irrational number}$$

$$\Rightarrow (2, \sqrt{3}) \in R$$

$$\text{Also } \sqrt{3} - \sqrt{2} + \sqrt{2} = \sqrt{3} \text{ which is an irrational number}$$

$$\Rightarrow (\sqrt{3}, \sqrt{2}) \in R$$

$$\text{But } 2 - \sqrt{2} + \sqrt{2} = 2 \text{ which is a rational number.}$$

$$\Rightarrow (2, \sqrt{2}) \notin R$$

$\Rightarrow R$ is not transitive

Option (a) is correct.

4. Given, $aRb \Rightarrow a$ is brother of b

This does not mean that b is also a brother of a because b can be a sister of a .

Hence, R is not symmetric.

$$\text{Again, } aRb \Rightarrow a \text{ is brother of } b \text{ and } bRc \Rightarrow b \text{ is brother of } c.$$

So, a is brother of c .

Hence, R is transitive.

Option (b) is correct.

5. We are given set $A = \{1, 2, 3\}$

Number of equivalence relation on A = number of possible portion of $\{1, 2, 3\}$

i.e., $3 = 1 + 1 + 1$ Only one combination

$3 = 1 + 2$ 3 Possible combination

$3 = 3$ 1 possible combination

i.e., (i) $\{\{1\}, \{2\}, \{3\}\}$ i.e., $\{(1, 1), (2, 2), (3, 3)\}$

(ii) $\{\{1, 2\}, \{3\}\}$ i.e., $\{(1, 2), (2, 1), (3, 3)\}$

(iii) $\{\{1, 3\}, \{2\}\}$ i.e., $\{(1, 3), (3, 1), (2, 2)\}$

(iv) $\{\{2, 3\}, \{1\}\}$ i.e., $\{(2, 3), (3, 2), (1, 1)\}$

(v) $\{\{1, 2, 3\}\}$ i.e., $\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)\}$

i.e., Total number of equivalence relation = 5

Option (d) is correct.

6. For $l, m \in L$

if $(l, m) \in R \Rightarrow l \perp m \Rightarrow m \perp l \Rightarrow (m, l) \in R$

$\therefore R$ is symmetric.

Option (b) is correct.

7. Required relation is reflexive and symmetric but not transitive is given by

$$R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1), (1, 3), (3, 1)\}$$

which is reflexive as $(a, a) \in R \forall a \in A$

which is symmetric as $(a, b) \in R \Rightarrow (b, a) \in R$ for $a, b \in A$

But $(2, 1), (1, 3) \in R \nRightarrow (2, 3) \in R$

Hence R is not transitive.

There is only one such relation.

Option (a) is correct.

9. We have $n(A) = m, n(B) = n$.

\therefore Number of relations defined from A to B

$$= \text{number of possible subsets of } A \times B = 2^{n(A \times B)} = 2^{mn}$$

Option (a) is correct.

10. The total number of injective mappings from the set containing n elements into the set containing m elements is ${}^m P_n$. So here it is ${}^4 P_3 = 4! = 24$.

Option (c) is correct.

11. We have $f: \mathbb{R} \rightarrow \mathbb{R}$: such that

$$f(x) = 2^x + 2^{|x|} = \begin{cases} 2^x + 2^x & \text{if } x \geq 0 \\ 2^x + 2^{-x} & \text{if } x < 0 \end{cases} = \begin{cases} 2^{x+1} & \text{if } x \geq 0 \\ 2^x + 2^{-x} & \text{if } x < 0 \end{cases}$$

$$\Rightarrow f'(x) = \begin{cases} 2^{x+1} \log 2 & \text{if } x \geq 0 \\ 2^x \log 2 + (-2^{-x} \log 2) & \text{if } x < 0 \end{cases} = \begin{cases} 2^{x+1} \log 2 & \text{if } x \geq 0 \\ \log 2 (2^x - 2^{-x}) & \text{if } x < 0 \end{cases}$$

$$\therefore f'(x) > 0 \forall x \geq 0 \text{ and } f'(x) < 0 \forall x < 0$$

$\Rightarrow f(x)$ is strictly increasing in $\mathbb{R} \Rightarrow f(x)$ is one-one.

Also, $f(x) \rightarrow \infty$ if $x \rightarrow \pm \infty$ and $f(x) > 0 \forall x \in \mathbb{R}$

$\Rightarrow f$ is an into function.

$\therefore f(x)$ is one-one and into function.

Option (c) is correct.

12. We have $n(A) = 5$ and $n(B) = 6$

\therefore Number of one-one mapping from A to $B = 6!$

As $n(A) < n(B)$

\Rightarrow There is no onto function from A to B . i.e., number of onto function = 0.

\therefore Number of one-one and onto functions from A to $B = 0$

Option (c) is correct.

13. $f: \mathbb{Z} \rightarrow \mathbb{Z}$ given by $f(x) = x + 2$

One-one For $x_1, x_2 \in \mathbb{Z}$ such that $x_1 \neq x_2 \Rightarrow x_1 + 2 \neq x_2 + 2$

$\Rightarrow f(x_1) \neq f(x_2)$

$\Rightarrow f$ is one-one.

Onto Let $y \in \mathbb{Z}$ (co-domain) such that

$$f(x) = y \Rightarrow x + 2 = y \Rightarrow x = y - 2$$

For $y \in \mathbb{Z}$ (co-domain), $\exists x = y - 2 \in \mathbb{Z}$ (domain) such that

$$f(x) = f(y - 2) = y - 2 + 2 = y$$

$\Rightarrow f$ is onto

As f is one-one and onto.

$\Rightarrow f$ is a bijective function.

Option (b) is correct.

14. Given that, $f(x) = x^2 - 4x + 5$

$$\text{Let } y = x^2 - 4x + 5$$

$$y = x^2 - 4x + 4 + 1 = (x - 2)^2 + 1$$

$$(x - 2)^2 = y - 1 \Rightarrow x - 2 = \sqrt{y - 1}$$

$$\Rightarrow x = 2 + \sqrt{y - 1}$$

$$\therefore y - 1 \geq 0, y \geq 1$$

$$\text{Range} = [1, \infty)$$

Option (b) is correct.

15. Let $f(x) = 17 \Rightarrow x^2 + 1 = 17$

$$\Rightarrow x = \pm 4 \Rightarrow \text{Pre image of } 17 \text{ are } \{4, -4\}$$

and let $f(x) = -3 \Rightarrow x^2 + 1 = -3 \Rightarrow x^2 = -4$ which is not true and hence -3 has no pre image

17. $\therefore b > 6$ and $a = b - 2$

$$\Rightarrow (6, 8) \in R \text{ as } 8 > 6 \text{ and } 6 = 8 - 2$$

Option (c) is correct.

18. f is not one-one because

$$f(-2) = (-2)^4 = 16$$

$$f(2) = (2)^4 = 16$$

i.e., -2 and $2 \in R$ (Domain) have same f -image in R (co-domain)

$\Rightarrow f$ is not one-one.

Also $f(x) = x^4$ never achieve negative value.

\Rightarrow All negative real number of co-domain R have no pre-image in Domain R .

$\Rightarrow f$ is not onto.

Hence, f is neither one-one nor onto.

Option (d) is correct.

19. f is one-one because

$$f(x_1) = f(x_2) \Rightarrow 3x_1 = 3x_2$$

$$\Rightarrow x_1 = x_2 \quad \forall x_1, x_2 \in R \quad (\text{Domain})$$

Also f is onto as

$$\text{Let } f(x) = y \Rightarrow 3x = y \Rightarrow x = \frac{y}{3}$$

$$\forall y \in R (\text{codomain}) \exists x = \frac{y}{3} \in R (\text{domain})$$

$$\text{such that } f(x) = f\left(\frac{y}{3}\right) = 3 \times \frac{y}{3} = y$$

$\Rightarrow f(x)$ is onto

Therefore, f is one-one onto.

Option (a) is correct.

21. Let x be the pre image of 5

$$\Rightarrow f(x) = 5$$

$$\Rightarrow x^2 + 1 = 5$$

$$\Rightarrow x^2 = 4$$

$$\Rightarrow x = \pm 2$$

i.e., pre-image of 5 is $-2, +2$.

Similarly if x be pre-image of -5

$$\Rightarrow f(x) = -5$$

$$\Rightarrow x^2 + 1 = -5$$

$$\Rightarrow x^2 = -6$$

$$x = \pm\sqrt{-6} \notin R$$

i.e., No real number is pre-image of -5 . Hence ϕ is the primage of -5 .

Option (c) is correct.

22. To find out the domain of f , we have to find out that value of x for which $f(x)$ is real.

$$\Rightarrow x^2 - 4 \geq 0$$

$$\Rightarrow (x+2)(x-2) \geq 0$$

$$\Rightarrow (x+2) \geq 0, (x-2) \geq 0 \text{ or } (x+2) \leq 0, (x-2) \leq 0$$

$$\Rightarrow x \geq -2, x \geq 2 \text{ or } x \leq -2, x \leq 2$$

$$\Rightarrow x \geq 2 \text{ or } x \leq -2$$

Domain of f is $(-\infty, -2] \cup [2, \infty)$

Option (c) is correct.

23. $f(-2) + f(0) + f(2) + f(5) = -2 + 0 + 4 + 15 = 17$

Option (b) is correct.

24. As R is reflexive relation on A , and for being reflexive $(a, a) \in R, \forall a \in A$

Therefore, the minimum number of ordered pair in R is n .

$$\Rightarrow m \geq n.$$

Option (a) is correct.

25. Given function is $f(x) = \log_{3+x}(x^2 - 1)$

It is obvious that $f(x)$ is defined when $x^2 - 1 > 0$, $3 + x > 0$ and $3 + x \neq 1$.

$$\text{Now, } x^2 - 1 > 0 \Rightarrow x^2 > 1$$

$$\Rightarrow x < -1 \text{ or } x > 1$$

$$3 + x > 0 \Rightarrow x > -3$$

$$3 + x \neq 1 \Rightarrow x \neq -2$$

Therefore, domain of the function $f(x) = (-3, -2) \cup (-2, -1) \cup (1, \infty)$

Option (c) is correct.

26. From definition of onto function,

$$\text{Range of function} = \text{Codomain of function} = [0, \frac{\pi}{2})$$

$$\Rightarrow 0 \leq \tan^{-1}(x^2 + x + a) < \frac{\pi}{2}$$

$$\rightarrow 0 \leq (x^2 + x + a) < \infty$$

$$\Rightarrow x^2 + x + a > 0 \quad \forall x \in R$$

Hence $D \leq 0$

$$\Rightarrow 1^2 - 4a \leq 0$$

$$\Rightarrow 4a \geq 1$$

$$\Rightarrow a \geq \frac{1}{4}$$

$$\Rightarrow a \in [\frac{1}{4}, \infty)$$

Option (b) is correct.

27. We have,

$f: R \rightarrow R$ and $g: R \rightarrow R$ are such that

$$f(x) = \begin{cases} 0, & x \in \text{rational} \\ x, & x \in \text{irrational} \end{cases}$$

$$\text{and } g(x) = \begin{cases} 0, & x \in \text{irrational} \\ x, & x \in \text{rational} \end{cases}$$

$\therefore (f - g): R \rightarrow R$ such that,

$$(f - g)(x) = \begin{cases} -x, & x \in \text{rational} \\ x, & x \in \text{irrational} \end{cases}$$

From definition of $(f - g): R \rightarrow R$, it is obvious that, each rational number of domain of $(f - g)(x)$, associate to its negative rational number in codomain/range and each irrational number of domain of $(f - g)(x)$, associate to same irrational number in codomain/range.

\Rightarrow For each $x \in \text{Domain of } (f - g)(x)$, there is only one value in codomain/range of $(f - g)(x)$.

Hence, $(f - g)(x)$ is one-one onto.

Option (a) is correct.

28. We have set $A = \{1, 2, 3, 4\}$ & relation

$$R = \{(1, 2), (3, 4)\} \text{ on } A$$

As for $(a, b) \in R, \exists (b, c) \in R$ such that
 $(a, c) \in R$.

Hence R is transitive.

So (b) is correct option.

Option (b) is correct.

29. $\because |A| = 4, |B| = 5$ so there does not exist.
one-one and onto $\because |B| > |A|$ so it is not onto.
So (a) is correct option.
Option (a) is correct.

30. Number of onto functions are given by
 $2^3 - {}^2C_1(2-1)^3 + {}^2C_2(2-2)^3$
 $= 8 - 2 \times 1 + 0 = 8 - 2 = 6$
Option (c) is correct.

31. Number of one-one function $= {}^6P_4$
 $= \frac{6!}{2!} = \frac{720}{2} = 360$
Option (c) is correct.

32. Number of one-one and onto function from A to B where $|A| = m$ is m .
 \therefore Number of one-one onto function $= \underline{4} = 24$
Option (b) is correct.

[**Note:** One-one onto function (bijective) from A to B is possible if A and B have same number of elements.]

34. Let R be a relation on the set of all integers \mathbb{Z} , defined by

$$aRb \Leftrightarrow a > b \quad \forall a, b \in \mathbb{Z}$$

(i) **Reflexive:** For $1 \in \mathbb{Z}$

$1 \not R 1$ as $1 \not > 1$ so $(1, 1) \notin R \Rightarrow R$ is not reflexive on \mathbb{Z}

(ii) **Symmetric:** $(3, 2) \in R$ as $3 > 2$

But $(2, 3) \notin R$ as $2 \not > 3$

Hence R is not symmetric on \mathbb{Z}

(iii) **Transitive:** Let $(a, b) \in R$ and $(b, c) \in R$ $a > b$ and $b > c$

Now $a > b > c \Rightarrow a > c \Rightarrow (a, c) \in R$

Hence R is a transitive relation on \mathbb{Z}

Option (c) is correct.

36. $f(x) = 4^x + 4^{|x|}$

One-one

Let $x_1, x_2 \in R$ (domain) such that

$$x_1 \neq x_2$$

$$\Rightarrow 4^{x_1} + 4^{|x_1|} \neq 4^{x_2} + 4^{|x_2|}$$

$$\Rightarrow f(x_1) \neq f(x_2)$$

f is one-one

Onto

For $0 \in R$ (Co-domain) there is no $x \in R$ (domain) such that $f(x) = 0$.

$\therefore f$ is not onto

Range of $f = R - \{0\} \subseteq R$

Hence f is one-one into function.

Option (a) is correct.

39. Here co-domain $= \left[0, \frac{\pi}{2}\right)$

For onto function, we have

$$\text{Co-domain} = \text{Range} = 0 \leq x < \frac{\pi}{2}$$

This is valid if $x^2 + x + 2a \geq 0$

[$\because f(x) \geq 0$ i.e. $Ax^2 + Bx + C \geq 0$ then $D \leq 0$ if $A > 0$.]

$$\text{i.e., } x^2 + x + 2a \geq 0 \Rightarrow 1^2 - 4 \times 1 \times 2a \leq 0$$

$$\rightarrow 1 - 8a \leq 0 \rightarrow 1 \leq 8a \rightarrow 8a \geq 1 \rightarrow a \geq \frac{1}{8}$$

$$\therefore a \in \left[\frac{1}{8}, \infty\right)$$

Option (d) is correct.

40. We are given that

$$(f(x))^2 - 4f(x)f'(x) + (f'(x))^2 = 0$$

$$\Rightarrow f'(x) = \frac{4f(x) \pm \sqrt{16(f(x))^2 - 4(f(x))^2}}{2}$$

$$\therefore f'(x) = \frac{4f(x) \pm 2f(x)\sqrt{4-1}}{2}$$

$$= \frac{4f(x) \pm 2\sqrt{3}f(x)}{2}$$

$$= f(x)(2 \pm \sqrt{3})$$

$$\Rightarrow \frac{f'(x)}{f(x)} = (2 \pm \sqrt{3})$$

Integrating, we get

$$\Rightarrow \log f(x) = (2 \pm \sqrt{3})x + C$$

$$\Rightarrow f(x) = e^{(2 \pm \sqrt{3})x + C} = e^C e^{(2 \pm \sqrt{3})x} = \lambda e^{(2 \pm \sqrt{3})x}$$

where $\lambda = e^C$

$$\Rightarrow f(x) = \lambda e^{(2 \pm \sqrt{3})x} = \lambda e^{(2 + \sqrt{3})x}, \lambda e^{(2 - \sqrt{3})x}$$

Option (c) is correct.

41. $f(x) = \tan^{-1}\left(\frac{2x}{1-x^2}\right) = 2 \tan^{-1}x$

$f(x)$ is one-one and onto

i.e., $f'(x) > 0$ or $f'(x) < 0$ and co-domain = range of $f(x)$

$$B = f(-1, 1) = (2 \tan^{-1}(-1), 2 \tan^{-1}(1))$$

$$= \left(2 \times \left(-\frac{\pi}{4} \right), 2 \times \frac{\pi}{4} \right) = \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$$

Option (c) is correct.

42. Let $f(x) = y = \frac{x}{1+|x|} \forall x \in \mathbb{R}, y \in \mathbb{R}$

$$\therefore f(x) = \frac{x}{1+x} \text{ or } \frac{x}{1-x} \text{ is one-one}$$

Here range of $f(x)$ is $\mathbb{R} - \{-1, 1\}$

But y can not have any of the values $-1, 1$ for some x .

$\therefore f(x)$ is not an onto function.

Option (c) is correct.

44. Number of onto function

$$\begin{aligned} &= 3^6 - {}^3C_1(3-1)^6 + {}^3C_2(3-2)^6 - {}^3C_3(3-3)^6 \\ &= 3^6 - 3 \times 2^6 + 3 \times 1 = 3^6 - 3 \times 2^6 + 3 \\ &= 3 \times (3^5 - 2^6 + 1) = 3(243 - 64 + 1) \\ &= 3 \times (244 - 64) = 3 \times 180 = 540 \end{aligned}$$

Option (d) is correct.

45. As $(7, 7) \notin R$, so R can not be reflexive

Again $(7, 8) \in R$ but $(8, 7) \notin R$, so R is not symmetric.

As $(7, 8), (8, 8) \in R \Rightarrow (7, 8) \in R \Rightarrow R$ is transitive.

Option (a) is correct.

46. Let $A = \{1, 2, 3, \dots, 12\}$, $n(A) = 12$ (say m)

$B = \{1, 2, 3, \dots, 11\}$, $n(B) = 11$ (say n)

\therefore Total number of function from A to $B = 11^{12}$

$$\begin{aligned} \therefore \text{Number of onto functions from } A \text{ to } B &= \sum_{r=1}^n (-1)^{n-r} {}^nC_r r^m \\ &= \text{coefficient of } x^m \text{ in } m! (e^x - 1)^n \dots (i) \end{aligned}$$

Putting $m = 12$, $n = 11$ and $r = 1, 2, 3, \dots, 11$. The number of onto functions is given by

$$\begin{aligned} &= (-1)^{11-1} {}^{11}C_1 1^{12} + (-1)^{11-2} {}^{11}C_2 2^{12} + (-1)^{11-3} {}^{11}C_3 3^{12} \\ &+ \dots + (-1)^{1-11} {}^{11}C_{10} 10^{12} + (-1)^{0-11} {}^{11}C_{11} 11^{12} \\ &= {}^{11}C_{11} 11^{12} - {}^{11}C_{10} 10^{12} + {}^{11}C_9 9^{12} + \dots + {}^{11}C_3 3^{12} - {}^{11}C_2 2^{12} + {}^{11}C_1 1^{12} \\ &= {}^{11}C_0 11^{12} - {}^{11}C_1 10^{12} + {}^{11}C_2 9^{12} + \dots + {}^{11}C_8 3^{12} - {}^{11}C_9 2^{12} + {}^{11}C_{10} 1^{12} \end{aligned}$$

Also, coefficient x^{12} in $12! (e^x - 1)^{11}$

$$= \text{coefficient of } x^{12} \text{ in } 12! \left(1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots + \infty - 1 \right)^{11}$$

$$= \text{coefficient of } x^{12} \text{ in } 12! \left(\frac{x}{1!} + \frac{x^2}{2!} + \dots + \infty \right)^{11}$$

$$= \text{coefficient of } x^{12} \text{ in } 12! x^{11} \left(\frac{1}{1!} + \frac{x}{2!} + \frac{x^2}{3!} + \dots + \infty \right)^{11}$$

$$= \text{coefficient of } x \text{ in } 12! \times \left[1 + \left(\frac{x}{2!} + \frac{x^2}{3!} + \dots + \infty \right) \right]^{11}$$

$$= \text{coefficient of } x \text{ in } 12! \left[{}^{11}C_0 1 + {}^{11}C_1 \left(\frac{x}{2!} + \frac{x^2}{3!} + \dots + \infty \right) + {}^{11}C_2 \left(\frac{x}{2!} + \frac{x^2}{3!} + \dots + \infty \right) + \dots \right]$$

$$= \text{coefficient of } x \text{ in } 12! \left[{}^{11}C_1 \left(\frac{1}{2!} \right) \right] = \frac{12! \times 11}{2!}$$

$$\text{Total number of functions which are not onto} = 11^{12} - \frac{12! \times 11}{2}$$

Option (d) is correct.

47. $\therefore f(x \cdot f(y)) = x^a y^b \quad \dots(i)$

Replacing x by $\frac{1}{f(y)}$, we have from (i)

$$f\left(x \cdot f(y)\right) = f\left(x \times \frac{1}{f(y)}\right) = \left(\frac{1}{f(y)}\right)^a y^b$$

$$f(1) = \frac{y^b}{(f(y))^a} \Rightarrow (f(y))^a = \frac{y^b}{f(1)}$$

$$\therefore \text{Put } y = 1, (f(1))^a = \frac{1^b}{f(1)} = \frac{1}{f(1)}$$

$$\Rightarrow (f(1))^{a+1} = 1$$

$$\Rightarrow f(1) = 1^{\left(\frac{1}{a+1}\right)} = 1$$

$$\Rightarrow f(1) = \frac{y^b}{(f(y))^a} = 1 \Rightarrow (f(y))^a = y^b$$

$$\Rightarrow f(y) = y^{b/a}$$

Replacing y as x , we have

$$f(x) = x^{b/a} \quad \dots(ii)$$

$$\therefore f(x \cdot y^{b/a}) = x^a y^b$$

$$\text{Let } y^{b/a} = t \Rightarrow y = t^{a/b}$$

$$f(x \cdot t) = x^a t^a \Rightarrow f(x) = x^a \quad \dots(iii)$$

Now from (ii) and (iii), we get

$$x^{b/a} = x^a \Rightarrow \frac{a}{b} = \frac{1}{a} \Rightarrow b = a^2$$

Option (c) is correct.

51. Number of relations that can be defined on $A = 2^{n^2}$

Option (c) is correct.

54. We have $R = \{(a, a), (b, b), (c, c), (b, c), (a, b)\}$

For $(b, c) \in R$, but $(c, b) \notin R$.

Hence R is not symmetric.

Also for $(a, b), (b, c) \in R$ but $(a, c) \notin R$.

$\Rightarrow R$ is not transitive.

As $(a, a) \in R \forall a \in A$

Hence R is reflexive.

Option (c) is correct.

55. Let $A = \{1, 2\}, B = \{a, b\}$

Let $R = \{(1, a), (1, b), (2, a), (2, b)\}$

Clearly R is a relation from A to B

But R is not a function.

As $(1, a), (1, b) \in R$ and $(2, a), (2, b) \in R$

Option (b) is correct.

56. $R = \{(1, 2)\}$, $A = \{1, 2, 3\}$

Clearly R is neither reflexive nor symmetric.

As $(1, 2) \in R$ but $\nexists (2, b) \in R$ for $b \in A$ such that $(1, b) \in R$.

Hence R is a transitive relation on A .

Option (b) is correct.

57. $R = \{(a, b) : a \geq b\}$

Reflexive

Clearly $(a, a) \in R \forall a \in R$.

Hence R is reflexive.

Symmetric

$\therefore (2, 1) \in R$ but $(1, 2) \notin R$

Hence R is not symmetric.

Transitive

Let (a, b) and $(b, c) \in R$

$\Rightarrow a \geq b$ and $b \geq c$

$\Rightarrow a \geq c$

Hence (a, b) and $(b, c) \in R \Rightarrow (a, c) \in R$

$\Rightarrow R$ is a transitive relation on R .

Option (b) is correct.

59. $R = \{(x, y) : |x^2 - y^2| < 7\}$

$R = \{(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (1, 2), (2, 3)\}$

Option (d) is correct.



STAND ALONE MCQs

(1 Mark each)

Q. 1. Let T be the set of all triangles in the Euclidean plane, and let a relation R on T be defined as aRb if a is congruent to $b \forall a, b \in T$. Then R is

- (A) reflexive but not transitive
- (B) transitive but not symmetric
- (C) equivalence relation
- (D) None of these

Ans. Option (C) is correct.

Explanation: Consider that aRb , if a is congruent to $b, \forall a, b \in T$.

Then, $aRa \Rightarrow a \cong a$,

Which is true for all $a \in T$

So, R is reflexive,

Let $aRb \Rightarrow a \cong b$

$\Rightarrow b \cong a$

$\Rightarrow bRa$

So, R is symmetric.

Let aRb and bRc

$\Rightarrow b \cong b$ and $b \cong a$

$\Rightarrow a \cong c \Rightarrow aRc$

So, R is transitive

Hence, R is equivalence relation.

Q. 2. Consider the non-empty set consisting of children in a family and a relation R defined as aRb if a is

brother of b . Then R is

- (A) symmetric but not transitive
- (B) transitive but not symmetric
- (C) neither symmetric nor transitive
- (D) both symmetric and transitive

Ans. Option (B) is correct.

Explanation: $aRb \Rightarrow a$ is brother of b .

This does not mean b is also a brother of a as b can be a sister of a .

Hence, R is not symmetric.

$aRb \Rightarrow a$ is brother of b

and $bRc \Rightarrow b$ is a brother of c .

So, a is brother of c .

Hence, R is transitive.

Q. 3. The maximum number of equivalence relations on the set $A = \{1, 2, 3\}$ are

- (A) 1
- (B) 2
- (C) 3
- (D) 5

Ans. Option (D) is correct.

Explanation: Given that, $A = \{1, 2, 3\}$

Now, number of equivalence relations are as follows:

$R_1 = \{(1, 1), (2, 2), (3, 3)\}$

$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$

$$R_3 = \{(1, 1), (2, 2), (3, 3), (1, 3), (3, 1)\}$$

$$R_4 = \{(1, 1), (2, 2), (3, 3), (2, 3), (3, 2)\}$$

$$R_5 = \{(1, 2, 3) \leftrightarrow A \times A = A^2\}$$

\therefore Maximum number of equivalence relations on the set $A = \{1, 2, 3\} = 5$

Q. 4. If a relation R on the set $\{1, 2, 3\}$ be defined by $R = \{(1, 2)\}$, then R is

- (A) reflexive (B) transitive
(C) symmetric (D) None of these

Ans. Option (B) is correct.

Explanation: R on the set $\{1, 2, 3\}$ is defined by $R = \{(1, 2)\}$

It is clear that R is transitive.

Q. 5. Let us define a relation R in R as aRb if $a \geq b$. Then R is

- (A) an equivalence relation
(B) reflexive, transitive but not symmetric
(C) symmetric, transitive but not reflexive
(D) neither transitive nor reflexive but symmetric.

Ans. Option (B) is correct.

Explanation: Given that, aRb if $a \geq b$

$\Rightarrow aRa \Rightarrow a \geq a$ which is true

Let aRb , $a \geq b$, then $b \geq a$ which is not true as R is not symmetric.

But aRb and bRc

$\Rightarrow a \geq b$ and $b \geq c$

$\Rightarrow a \geq c$

Hence, R is transitive.

Q. 6. Let $A = \{1, 2, 3\}$ and consider the relation $R = (1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)$.

Then R is

- (A) reflexive but not symmetric
(B) reflexive but not transitive
(C) symmetric and transitive
(D) neither symmetric, nor transitive

Ans. Option (A) is correct.

Explanation: Given that $A = \{1, 2, 3\}$ and $R = (1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)$.

$\therefore (1, 1), (2, 2), (3, 3) \in R$

Hence, R is reflexive.

$(1, 2) \in R$ but $(2, 1) \notin R$

Hence, R is not symmetric.

$(1, 2) \in R$ and $(2, 3) \in R$

$\Rightarrow (1, 3) \in R$

Hence, R is transitive.

Q. 7. Let R be the relation in the set $\{1, 2, 3, 4\}$ given by $R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$. Choose the correct answer:

- (A) R is reflexive and symmetric but not transitive
(B) R is reflexive and transitive but not symmetric
(C) R is symmetric and transitive but not reflexive
(D) R is an equivalence relation

Ans. Option (B) is correct.

Explanation: Let R be the relation in the set $\{1, 2, 3, 4\}$ is given by:

$$R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$$

(a) $(1, 1), (2, 2), (3, 3), (4, 4) \in R$

Therefore, R is reflexive.

(b) $(1, 2) \in R$ but $(2, 1) \notin R$.

Therefore, R is not symmetric.

(c) If $(1, 3) \in R$ and $(3, 2) \in R$ then $(1, 2) \in R$.

Therefore, R is transitive.

Q. 8. Let $A = \{1, 2, 3\}$. Then number of relations containing $(1, 2)$ and $(1, 3)$ which are reflexive and symmetric but not transitive is

- (A) 1 (B) 2
(C) 3 (D) 4

Ans. Option (A) is correct.

Explanation: The given set is $A = \{1, 2, 3\}$.

The smallest relation containing $(1, 2)$ and $(1, 3)$, which is reflexive and symmetric, but not transitive is given by:

$$R = \{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 1), (3, 1)\}$$

This is because relation R is reflexive as

$$(1, 1), (2, 2), (3, 3) \in R.$$

Relation R is symmetric since $(1, 2), (2, 1) \in R$ and $(1, 3), (3, 1) \in R$.

But relation R is not transitive as $(3, 1), (1, 2) \in R$, but $(3, 2) \notin R$.

Now, if we add any two pairs $(3, 2)$ and $(2, 3)$ (or both) to relation R , then relation R will become transitive.

Hence, the total number of desired relations is one.

Q. 9. If the set A contains 5 elements and the set B contains 6 elements, then the number of one-one and onto mappings from A to B is

- (A) 720 (B) 120
(C) 0 (D) None of these

Ans. Option (C) is correct.

Explanation: We know that, if A and B are two non-empty finite sets containing m and n elements, respectively, then the number of one-one and onto mapping from A to B is

$n!$ if $m = n$

0, if $m \neq n$

Given that, $m = 5$ and $n = 6$

$\therefore m \neq n$

Number of one-one and onto mapping = 0

Q. 10. Let $A = \{1, 2, 3, \dots, n\}$ and $B = \{a, b\}$. Then the number of surjections from A into B is

- (A) nP_2 (B) $2^n - 2$
(C) $2^n - 1$ (D) None of these

Ans. Option (B) is correct.

Explanation: Total number of functions from A to $B = 2^n$

Number of into functions = 2

Number of surjections from A to $B = 2^n - 2$

Q. 11. Let $f: R \rightarrow R$ be defined by $f(x) = \frac{1}{x}, \forall x \in R$. Then f is

- (A) one-one (B) onto
(C) bijective (D) f is not defined

Ans. Option (D) is correct.

Explanation: We have, $f(x) = \frac{1}{x}, \forall x \in R$

For $x = 0$, $f(x)$ is not defined.

Hence, $f(x)$ is a not defined function.

Q. 12. Which of the following functions from Z into Z are bijections?

- (A) $f(x) = x^3$ (B) $f(x) = x + 2$
(C) $f(x) = 2x + 1$ (D) $f(x) = x^2 + 1$

Ans. Option (B) is correct.

Explanation: For bijection on Z , $f(x)$ must be one-one and onto.

Function $f(x) = x^2 + 1$ is many-one as $f(1) = f(-1)$

Range of $f(x) = x^3$ is not Z for $x \in Z$.

Also $f(x) = 2x + 1$ takes only values of type $= 2k + 1$ for $x \in k \in Z$

But $f(x) = x + 2$ takes all integral values for $x \in Z$

Hence $f(x) = x + 2$ is bijection of Z .

Q. 13. Let $f: R \rightarrow R$ be defined as $f(x) = x^4$. Choose the correct answer.

- (A) f is one-one onto
(B) f is many-one onto
(C) f is one-one but not onto

(D) f is neither one-one nor onto

Ans. Option (D) is correct.

Explanation: We know that $f: R \rightarrow R$ is defined as $f(x) = x^4$.

Let $x, y \in R$ such that $f(x) = f(y)$

$$\Rightarrow x^4 = y^4$$

$$\Rightarrow x = \pm y$$

$$\therefore f(x) = f(y)$$

does not imply that $x = y$.

For example, $f(1) = f(-1) = 1$

$\therefore f$ is not one-one.

Consider an element 2 in co-domain R . It is clear that there does not exist any x in domain R such that $f(x) = 2$.

$\therefore f$ is not onto.

Hence, function f is neither one-one nor onto.

Q. 14. Let $f: R \rightarrow R$ be defined as $f(x) = 3x$. Choose the correct answer.

- (A) f is one-one onto
(B) f is many-one onto
(C) f is one-one but not onto
(D) f is neither one-one nor onto

Ans. Option (A) is correct.

Explanation: $f: R \rightarrow R$ is defined as $f(x) = 3x$.

Let $x, y \in R$ such that $f(x) = f(y)$

$$\Rightarrow 3x = 3y$$

$$\Rightarrow x = y$$

$\therefore f$ is one-one.

Also, for any real number y in co-domain R , there

exists $\frac{y}{3}$ in R such that $f\left(\frac{y}{3}\right) = 3\left(\frac{y}{3}\right) = y$.

$\therefore f$ is onto.

Hence, function f is one-one and onto.



ASSERTION AND REASON BASED MCQs

(1 Mark each)

Directions: In the following questions, A statement of Assertion (A) is followed by a statement of Reason (R). Mark the correct choice as.

- (A) Both A and R are true and R is the correct explanation of A
(B) Both A and R are true but R is NOT the correct explanation of A
(C) A is true but R is false
(D) A is false and R is True

Q. 1. Let W be the set of words in the English dictionary. A relation R is defined on W as

$R = \{(x, y) \in W \times W \text{ such that } x \text{ and } y \text{ have at least one letter in common}\}$.

Assertion (A): R is reflexive.

Reason (R): R is symmetric.

Ans. Option (B) is correct.

Explanation: For any word $x \in W$

x and x have atleast one (all) letter in common

$\therefore (x, x) \in R, \forall x \in W \therefore R$ is reflexive

Symmetric: Let $(x, y) \in R, x, y \in W$

$\Rightarrow x$ and y have atleast one letter in common

$\Rightarrow y$ and x have atleast one letter in common

$\Rightarrow (y, x) \in R \therefore R$ is symmetric

Hence A is true, R is true; R is not a correct explanation for A.

Q. 2. Let R be the relation in the set of integers Z given by $R = \{(a, b) : 2 \text{ divides } a - b\}$.

Assertion (A): R is a reflexive relation.

Reason (R): A relation is said to be reflexive if $xRx, \forall x \in Z$.

Ans. Option (A) is correct.

Explanation: By definition, a relation in Z is said to be reflexive if $xRx, \forall x \in Z$. So R is true.

$$a - a = 0 \Rightarrow 2 \text{ divides } a - a \Rightarrow aRa.$$

Hence R is reflexive and A is true.

R is the correct explanation for A .

Q. 3. Consider the set $A = \{1, 3, 5\}$.

Assertion (A): The number of reflexive relations on set A is 2^9 .

Reason (R): A relation is said to be reflexive if $xRx, \forall x \in A$.

Ans. Option (D) is correct.

Explanation: By definition, a relation in A is said to be reflexive if $xRx, \forall x \in A$. So R is true.

The number of reflexive relations on a set containing n elements is 2^{n^2-n} .

Here $n = 3$.

The number of reflexive relations on a set $A = 2^{9-3} = 2^6$.

Hence A is false.

Q. 4. Consider the function $f: R \rightarrow R$ defined as $f(x) = x^3$

Assertion (A): $f(x)$ is a one-one function.

Reason (R): $f(x)$ is a one-one function if co-domain = range.

Ans. Option (C) is correct.

Explanation: $f(x)$ is a one-one function if

$$f(x_1) = f(x_2) \Rightarrow x_1 = x_2.$$

Hence R is false.

Let $f(x_1) = f(x_2)$ for some $x_1, x_2 \in R$

$$\Rightarrow (x_1)^3 = (x_2)^3$$

$$\Rightarrow x_1 = x_2$$

Hence $f(x)$ is one-one.

Hence A is true.

Q. 5. If $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and $f = \{(1, 4), (2, 5), (3, 6)\}$ is a function from A to B .

Assertion (A): $f(x)$ is a one-one function.

Reason (R): $f(x)$ is an onto function.

Ans. Option (C) is correct.

Given, $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and $f: A \rightarrow B$ is defined as $f = \{(1, 4), (2, 5), (3, 6)\}$ i.e., $f(1) = 4$, $f(2) = 5$ and $f(3) = 6$.

It can be seen that the images of distinct elements of A under f are distinct. So, f is one-one.

So, A is true.

Range of $f = \{4, 5, 6\}$.

Co-domain = $\{4, 5, 6, 7\}$.

Since co-domain \neq range, $f(x)$ is not an onto function. Hence R is false.

Q. 6. Consider the function $f: R \rightarrow R$ defined as

$$f(x) = \frac{x}{x^2 + 1}.$$

Assertion (A): $f(x)$ is not one-one.

Reason (R): $f(x)$ is not onto.

Ans. Option (B) is correct.

Explanation: Given, $f: R \rightarrow R$;

$$f(x) = \frac{x}{1+x^2}$$

$$\text{Taking } x_1 = 4, x_2 = \frac{1}{4} \in R$$

$$f(x_1) = f(4) = \frac{4}{17}$$

$$f(x_2) = f\left(\frac{1}{4}\right) = \frac{4}{17} \quad (x_1 \neq x_2)$$

$\therefore f$ is not one-one.

A is true.

Let $y \in R$ (co-domain)

$$f(x) = y$$

$$\Rightarrow \frac{x}{1+x^2} = y$$

$$\Rightarrow y(1+x^2) = x$$

$$\Rightarrow yx^2 + y - x = 0$$

$$\Rightarrow x = \frac{1 \pm \sqrt{1-4y^2}}{2y}$$

since, $x \in R$,

$$\therefore 1 - 4y^2 \geq 0$$

$$\Rightarrow -\frac{1}{2} \leq y \leq \frac{1}{2}$$

$$\text{So Range } (f) \in \left[-\frac{1}{2}, \frac{1}{2}\right]$$

Range $(f) \neq R$ (Co-domain)

$\therefore f$ is not onto.

R is true.

R is not the correct explanation for A .



CASE-BASED MCQs

Attempt any four sub-parts from each question. Each sub-part carries 1 mark.

I. Read the following text and answer the following questions on the basis of the same:

A general election of Lok Sabha is a gigantic exercise.

About 911 million people were eligible to vote and voter turnout was about 67%, the highest ever

Let I be the set of all citizens of India who were eligible to exercise their voting right in general election held in 2019. A relation ' R ' is defined on I as follows:

ONE – NATION
ONE – ELECTION
FESTIVAL OF DEMOCRACY
GENERAL ELECTION - 2019



$R = \{(V_1, V_2) : V_1, V_2 \in I \text{ and both use their voting right in general election - 2019}\}$

[CBSE QB 2021]

Q. 1. Two neighbours X and $Y \in I$. X exercised his voting right while Y did not cast her vote in general election - 2019. Which of the following is true?

- (A) $(X, Y) \in R$ (B) $(Y, X) \in R$
(C) $(X, X) \notin R$ (D) $(X, Y) \notin R$

Ans. Option (D) is correct.

Explanation: $(X, Y) \notin R$.

$\because X$ exercised his voting right while, Y did not cast her vote in general election-2019

And $R = \{(V_1, V_2) : V_1, V_2 \in I \text{ and both use their voting right in general election-2019}\}$

Q. 2. Mr. 'X' and his wife 'W' both exercised their voting right in general election -2019, Which of the following is true?

- (A) both (X, W) and $(W, X) \in R$
(B) $(X, W) \in R$ but $(W, X) \notin R$
(C) both (X, W) and $(W, X) \notin R$
(D) $(W, X) \in R$ but $(X, W) \notin R$

Ans. Option (A) is correct.

Q. 3. Three friends F_1, F_2 and F_3 exercised their voting right in general election-2019, then which of the following is true?

- (A) $(F_1, F_2) \in R, (F_2, F_3) \in R$ and $(F_1, F_3) \in R$
(B) $(F_1, F_2) \in R, (F_2, F_3) \in R$ and $(F_1, F_3) \notin R$
(C) $(F_1, F_2) \in R, (F_2, F_3) \in R$ but $(F_1, F_3) \notin R$
(D) $(F_1, F_2) \notin R, (F_2, F_3) \notin R$ and $(F_1, F_3) \in R$

Ans. Option (A) is correct.

Q. 4. The above defined relation R is _____

- (A) Symmetric and transitive but not reflexive
(B) Universal relation
(C) Equivalence relation
(D) Reflexive but not symmetric and transitive

Ans. Option (C) is correct.

Explanation: R is reflexive, since every person is friend or itself.

i.e., $(F_1, F_2) \in R$

Further, $(F_1, F_2) \in R$

$\Rightarrow F_1$ is friend of F_2

$\Rightarrow F_2$ is friend of F_1

$\Rightarrow (F_2, F_1) \in R$

$\Rightarrow R$ is symmetric

Moreover, $(F_1, F_2), (F_2, F_3) \in R$

$\Rightarrow F_1$ is friend of F_2 and F_2 is friend of F_3 .

$\Rightarrow F_1$ is a friend of F_3 .

$\Rightarrow (F_1, F_3) \in R$

Therefore, R is an equivalence relation.

Q. 5. Mr. Shyam exercised his voting right in General Election - 2019, then Mr. Shyam is related to which of the following?

- (A) All those eligible voters who cast their votes
(B) Family members of Mr. Shyam
(C) All citizens of India
(D) Eligible voters of India

Ans. Option (A) is correct.

II. Read the following text and answer the following questions on the basis of the same:

Sherlin and Danju are playing Ludo at home during Covid-19. While rolling the dice, Sherlin's sister Raji observed and noted the possible outcomes of the throw every time belongs to set $\{1, 2, 3, 4, 5, 6\}$. Let A be the set of players while B be the set of all possible outcomes.



$A = \{S, D\}, B = \{1, 2, 3, 4, 5, 6\}$ [CBSE QB 2021]

Q. 1. Let $R : B \rightarrow B$ be defined by $R = \{(x, y) : y \text{ is divisible by } x\}$ is

- (A) Reflexive and transitive but not symmetric
(B) Reflexive and symmetric but not transitive
(C) Not reflexive but symmetric and transitive
(D) Equivalence

Ans. Option (A) is correct.

Explanation: R is reflexive, since every element of B i.e.,

$B = \{1, 2, 3, 4, 5, 6\}$ is divisible by itself.

i.e., $(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6) \in R$

further, $(1, 2) \in R$

but $(2, 1) \notin R$

Moreover,

$(1, 2), (2, 4) \in R$

$\Rightarrow (1, 4) \in R$

$\Rightarrow R$ is transitive.

Therefore, R is reflexive and transitive but not symmetric.

Q. 2. Raji wants to know the number of functions from A to B . How many number of functions are possible?

- (A) 6^2 (B) 2^6
(C) $6!$ (D) 2^{12}

Ans. Option (A) is correct.

- Q. 3. Let R be a relation on B defined by $R = \{(1, 2), (2, 2), (1, 3), (3, 4), (3, 1), (4, 3), (5, 5)\}$. Then R is
(A) Symmetric
(B) Reflexive
(C) Transitive
(D) None of these

Ans. Option (D) is correct.

Explanation: $R = \{(1, 2), (2, 2), (1, 3), (3, 4), (3, 1), (4, 3), (5, 5)\}$
 R is not reflexive.
Since, $(1, 1), (3, 3), (4, 4), (6, 6) \in R$
 R is not symmetric.
Because, for $(1, 2) \in R$ there does not exist $(2, 1) \in R$.
 R is not transitive.
Because for all element of B there does not exist,
 $(a, b) (b, c) \in R$ and $(a, c) \in R$.

- Q. 4. Raji wants to know the number of relations possible from A to B . How many numbers of relations are possible?
(A) 6^2 (B) 2^6
(C) $6!$ (D) 2^{12}

Ans. Option (D) is correct.

- Q. 5. Let $R : B \rightarrow B$ be defined by $R = \{(1, 1), (1, 2), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}$, then R is
(A) Symmetric
(B) Reflexive and Transitive
(C) Transitive and symmetric
(D) Equivalence

Ans. Option (B) is correct.

III. Read the following text and answer the following questions on the basis of the same:

An organization conducted bike race under 2 different categories—boys and girls. Totally there were 250 participants. Among all of them finally three from Category 1 and two from Category 2 were selected for the final race. Ravi forms two sets B and G with these participants for his college project.

Let $B = \{b_1, b_2, b_3\}$ $G = \{g_1, g_2\}$ where B represents the set of boys selected and G the set of girls who were selected for the final race. [CBSE QB 2021]



Ravi decides to explore these sets for various types of relations and functions

- Q. 1. Ravi wishes to form all the relations possible from B to G . How many such relations are possible?
(A) 2^6 (B) 2^5
(C) 0 (D) 2^3

Ans. Option (A) is correct.

- Q. 2. Let $R : B \rightarrow B$ be defined by $R = \{(x, y) : x \text{ and } y \text{ are students of same sex}\}$, Then this relation R is _____
(A) Equivalence
(B) Reflexive only
(C) Reflexive and symmetric but not transitive
(D) Reflexive and transitive but not symmetric

Ans. Option (A) is correct.

Explanation:

$R : B \rightarrow B$ be defined by $R = \{(x, y) : x \text{ and } y \text{ are students of same sex}\}$
 R is reflexive, since, $(x, x) \in R$
 R is symmetric, since, $(x, y) \in R$ and $(y, x) \in R$
 R is transitive. For $a, b, c \in B$
 $\exists (a, b) (b, c) \in R$
and $(a, c) \in R$.
Therefore R is equivalence relation.

- Q. 3. Ravi wants to know among those relations, how many functions can be formed from B to G ?
(A) 2^2 (B) 2^{12}
(C) 3^2 (D) 2^3

Ans. Option (D) is correct.

- Q. 4. Let $R : B \rightarrow G$ be defined by $R = \{(b_1, g_1), (b_2, g_2), (b_3, g_1)\}$, then R is _____
(A) Injective
(B) Surjective
(C) Neither Surjective nor Injective
(D) Surjective and Injective

Ans. Option (B) is correct.

Explanation:

$R : B \rightarrow G$ be defined by $R = \{(b_1, g_1), (b_2, g_2), (b_3, g_1)\}$
 R is surjective, since, every element of G is the image of some element of B under R , i.e., For $g_1, g_2 \in G$,
there exists an elements $b_1, b_2, b_3 \in B$,
 $(b_1, g_1) (b_2, g_2), (b_3, g_1) \in R$.

- Q. 5. Ravi wants to find the number of injective functions from B to G . How many numbers of injective functions are possible?
(A) 0 (B) $2!$
(C) $3!$ (D) $0!$

Ans. Option (A) is correct.

IV. Read the following text and answer the following questions on the basis of the same:

Students of Grade 9, planned to plant saplings

along straight lines, parallel to each other to one side of the playground ensuring that they had enough play area. Let us assume that they planted one of the rows of the saplings along the line $y = x - 4$. Let L be the set of all lines which are parallel on the ground and R be a relation on L .

[CBSE QB 2021]



Q. 1. Let relation R be defined by $R = \{(L_1, L_2) : L_1 \parallel L_2 \text{ where } L_1, L_2 \in L\}$ then R is _____ relation

- (A) Equivalence
- (B) Only reflexive
- (C) Not reflexive
- (D) Symmetric but not transitive

Ans. Option (A) is correct.

Explanation: Let relation R be defined by $R = \{(L_1, L_2) : L_1 \parallel L_2 \text{ where } L_1, L_2 \in L\}$.
 R is reflexive, since every line is parallel to itself.
Further, $(L_1, L_2) \in R$
 $\Rightarrow L_1$ is parallel to L_2
 $\Rightarrow L_2$ is parallel to L_1
 $\Rightarrow (L_2, L_1) \in R$
Hence, R is symmetric.
Moreover, $(L_1, L_2), (L_2, L_3) \in R$
 $\Rightarrow L_1$ is parallel to L_2 and L_2 is parallel to L_3
 $\Rightarrow L_1$ is parallel to L_3
 $\Rightarrow (L_1, L_3) \in R$
Therefore, R is an equivalence relation

Q. 2. Let $R = \{(L_1, L_2) : L_1 \perp L_2 \text{ where } L_1, L_2 \in L\}$ which of the following is true?

- (A) R is Symmetric but neither reflexive nor transitive
- (B) R is Reflexive and transitive but not symmetric
- (C) R is Reflexive but neither symmetric nor transitive
- (D) R is an Equivalence relation

Ans. Option (A) is correct.

Explanation: R is not reflexive, as a line L_1 can not be perpendicular to itself, i.e., $(L_1, L_1) \notin R$.
 R is symmetric as $(L_1, L_2) \in R$
As, L_1 is perpendicular to L_2

and L_2 is perpendicular to L_1
 $(L_2, L_1) \in R$

R is not transitive. Indeed, if L_1 is perpendicular to L_2 and L_2 is perpendicular to L_3 , then L_1 can never be perpendicular to L_3 .

In fact L_1 is parallel to L_3 ,
i.e., $(L_1, L_2) \in R, (L_2, L_3) \in R$ but $(L_1, L_3) \notin R$
i.e., symmetric but neither reflexive nor transitive.

Q. 3. The function $f : R \rightarrow R$ defined by $f(x) = x - 4$ is _____

- (A) Bijective
- (B) Surjective but not injective
- (C) Injective but not Surjective
- (D) Neither Surjective nor Injective

Ans. Option (A) is correct.

Explanation:

The function f is one-one,
for $f(x_1) = f(x_2)$
 $\Rightarrow x_1 - 4 = x_2 - 4$
 $\Rightarrow x_1 = x_2$

Also, given any real number y in R , there exists $y + 4$ in R

Such that $f(y + 4) = y + 4 - 4 = y$

Hence, f is onto

Hence, function is both one-one and onto, i.e., bijective.

Q. 4. Let $f : R \rightarrow R$ be defined by $f(x) = x - 4$. Then the range of $f(x)$ is _____

- (A) R
- (B) Z
- (C) W
- (D) Q

Ans. Option (A) is correct.

Explanation: Range of $f(x)$ is R

Q. 5. Let $R = \{(L_1, L_2) : L_1 \parallel L_2 \text{ and } L_1 : y = x - 4\}$ then which of the following can be taken as L_2 ?

- (A) $2x - 2y + 5 = 0$
- (B) $2x + y = 5$
- (C) $2x + 2y + 7 = 0$
- (D) $x + y = 7$

Ans. Option (A) is correct.

Explanation: Since, $L_1 \parallel L_2$
then slope of both the lines should be same.

Slope of $L_1 = 1$
 \Rightarrow Slope of $L_2 = 1$
And $2x - 2y + 5 = 0$
 $-2y = -2x - 5$
 $y = x + \frac{5}{2}$

Slope of $2x - 2y + 5 = 0$ is 1

So, $2x - 2y + 5 = 0$ can be taken as L_2 .

V. Read the following text and answer the following questions on the basis of the same:

Raji visited the Exhibition along with her family. The Exhibition had a huge swing, which attracted

many children. Raji found that the swing traced the path of a Parabola as given by $y = x^2$.

[CBSE QB-2021]



Q. 1. Let $f: R \rightarrow R$ be defined by $f(x) = x^2$ is _____

- (A) Neither Surjective nor Injective
- (B) Surjective
- (C) Injective
- (D) Bijective

Ans. Option (A) is correct.

Explanation:

$f: R \rightarrow R$ be defined by $f(x) = x^2$

$\therefore f(-1) = f(1) = 1$, but $-1 \neq 1$

$\therefore f$ is not injective

Now, $-2 \in R$. But, there does not exist any element $x \in R$ such that $f(x) = -2$ or $x^2 = -2$

$\therefore f$ is not surjective.

Hence, function f is neither injective nor surjective.

Q. 2. Let $f: N \rightarrow N$ be defined by $f(x) = x^2$ is _____

- (A) Surjective but not Injective
- (B) Surjective
- (C) Injective
- (D) Bijective

Ans. Option (C) is correct.

Explanation: $f: N \rightarrow N$ be defined by $f(x) = x^2$

for $x, y \in N$, $f(x) = f(y)$

$\Rightarrow x^2 = y^2$

$\Rightarrow x = y$

$\therefore f$ is injective

Now, $2 \in N$, But, there does not exist any x in n such that $f(x) = x^2 = 2$

$\therefore f$ is not surjective

Hence, function is injective but not surjective.

Q. 3. Let $f: \{1, 2, 3, \dots\} \rightarrow \{1, 4, 9, \dots\}$ be defined by $f(x) = x^2$ is _____

- (A) Bijective
- (B) Surjective but not Injective
- (C) Injective but Surjective
- (D) Neither Surjective nor Injective

Ans. Option (A) is correct.

Explanation:

$f: \{1, 2, 3, \dots\} \rightarrow \{1, 4, 9, \dots\}$ be defined by

$f(x) = x^2$

$x_1 \in \{1, 2, 3, \dots\}$ and $x_2 \in \{1, 2, 3, \dots\}$

$f(x_1) = f(x_2)$

$\Rightarrow x_1^2 = x_2^2$

$\Rightarrow x_1 = x_2$

$\therefore f$ is injective

Now, $4 \in \{1, 4, 9, \dots\}$, there exist 2 in $\{1, 2, 3, \dots\}$ such that $f(x) = 2^2 = 4$, Hence, f is surjective

Therefore f is bijective.

Q. 4. Let $f: N \rightarrow R$ be defined by $f(x) = x^2$. Range of the function among the following is _____

- (A) $\{1, 4, 9, 16, \dots\}$
- (B) $\{1, 4, 8, 9, 10, \dots\}$
- (C) $\{1, 4, 9, 15, 16, \dots\}$
- (D) $\{1, 4, 8, 16, \dots\}$

Ans. Option (A) is correct.

Explanation:

Range of $f = \{1, 4, 9, 16, \dots\}$

$\therefore N = \{1, 2, 3, \dots\}$

Q. 5. The function $f: Z \rightarrow Z$ defined by $f(x) = x^2$ is _____

- (A) Neither Injective nor Surjective
- (B) Injective
- (C) Surjective
- (D) Bijective

Ans. Option (A) is correct.

Explanation: $f: z \rightarrow z$ defined by $f(x) = x^2$

So, $f(-1) = f(1)$, but $-1 \neq 1$

$\therefore f$ is not injective

Now, $-2 \in Z$, but, there does not exist any element $x \in z$ such that

$f(x) = -2$

or $x^2 = -2$

$\therefore f$ is not surjective

Hence, f is neither injective nor surjective.