



www.aepstudycircle.com



CBSE
MATHEMATICS

+91-9939586130
+91-7739650505






QUICK RECAP

VECTOR

▶▶ A physical quantity having magnitude as well as direction is called a vector. A vector is represented by a line segment, denoted as \overline{AB} or \vec{a} . Here, point A is the initial point and B is the terminal point of the vector \overline{AB} .

- ▶▶ **Magnitude** : The distance between the points A and B is called the magnitude of the directed line segment \overline{AB} . It is denoted by $|\overline{AB}|$.
- ▶▶ **Position Vector** : Let P be any point in space, having coordinates (x, y, z) with respect to some fixed point $O(0, 0, 0)$ as origin, then

the vector \overline{OP} having O as its initial point and P as its terminal point is called the position vector of the point P with respect to O . The vector \overline{OP} is usually denoted by \vec{r} .



Magnitude of \overline{OP} is, $|\overline{OP}| = \sqrt{x^2 + y^2 + z^2}$
 i.e., $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$.

In general, the position vectors of points A , B , C , etc. with respect to the origin O are denoted by \vec{a} , \vec{b} , \vec{c} , etc. respectively.

▶▶ **Direction Cosines and Direction Ratios :**
 The angles α , β , γ made by the vector \vec{r} with the positive directions of x , y and z -axes respectively are called its direction angles. The cosine values of these angles, i.e., $\cos\alpha$, $\cos\beta$ and $\cos\gamma$ are called direction cosines of the vector \vec{r} , and usually denoted by l , m and n respectively.

Direction cosines of \vec{r} are given as

$$l = \frac{x}{\sqrt{x^2 + y^2 + z^2}}, m = \frac{y}{\sqrt{x^2 + y^2 + z^2}}, n = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$

The numbers lr , mr and nr , proportional to the direction cosines of vector \vec{r} are called direction ratios of the vector \vec{r} and denoted as a , b and c respectively.

i.e., $a = lr$, $b = mr$ and $c = nr$

Note : $l^2 + m^2 + n^2 = 1$ and $a^2 + b^2 + c^2 \neq 1$, (in general).

TYPES OF VECTORS


- ▶▶ **Zero vector :** A vector whose initial and terminal points coincide is called a zero (or null) vector. It cannot be assigned a definite direction as it has zero magnitude and it is denoted by the $\vec{0}$.
- ▶▶ **Unit Vector :** A vector whose magnitude is unity i.e., $|\vec{a}| = 1$. It is denoted by \hat{a} .
- ▶▶ **Equal Vectors :** Two vectors \vec{a} and \vec{b} are said to be equal, written as $\vec{a} = \vec{b}$, iff

they have equal magnitudes and direction regardless of the positions of their initial points.


- ▶▶ **Coinitial Vectors :** Vectors having same initial point are called co-initial vectors.
- ▶▶ **Collinear Vectors :** Two or more vectors are called collinear if they have same or parallel supports, irrespective of their magnitudes and directions.
- ▶▶ **Negative of a Vector :** A vector having the same magnitude as that of a given vector but directed in the opposite sense is called negative of the given vector i.e., $\overline{BA} = -\overline{AB}$.

ADDITION OF VECTORS

▶▶ **Triangle law :** Let the vectors \vec{a} and \vec{b} so positioned such that initial point of one coincides with terminal point of the other. If $\vec{a} = \overline{AB}$, $\vec{b} = \overline{BC}$. Then the vector $\vec{a} + \vec{b}$ is represented by the third side of $\triangle ABC$ i.e., $\overline{AB} + \overline{BC} = \overline{AC}$



▶▶ **Parallelogram law :** If the two vectors \vec{a} and \vec{b} are represented by the two adjacent sides OA and OB of a parallelogram $OACB$, then their sum $\vec{a} + \vec{b}$ is represented in magnitude and direction by the diagonal OC of parallelogram $OACB$ through their common point O i.e., $\overline{OA} + \overline{OB} = \overline{OC}$



Properties of Vector Addition

- ▶ Vector addition is commutative i.e., $\vec{a} + \vec{b} = \vec{b} + \vec{a}$.
- ▶ Vector addition is associative i.e., $\vec{a} + (\vec{b} + \vec{c}) = (\vec{a} + \vec{b}) + \vec{c}$.
- ▶ Existence of additive identity : The zero vector acts as additive identity i.e., $\vec{a} + \vec{0} = \vec{a} = \vec{0} + \vec{a}$ for any vector \vec{a} .
- ▶ Existence of additive inverse : The negative of \vec{a} i.e., $-\vec{a}$ acts as additive inverse i.e., $\vec{a} + (-\vec{a}) = \vec{0} = (-\vec{a}) + \vec{a}$ for any vector \vec{a} .

MULTIPLICATION OF A VECTOR BY A SCALAR

▶ Let \vec{a} be a given vector and λ be a given scalar (a real number), then $\lambda\vec{a}$ is defined as the multiplication of vector \vec{a} by the scalar λ . Its magnitude is $|\lambda|$ times the modulus of \vec{a} i.e., $|\lambda\vec{a}| = |\lambda| |\vec{a}|$.

Direction of $\lambda\vec{a}$ is same as that of \vec{a} if $\lambda > 0$ and opposite to that of \vec{a} if $\lambda < 0$.

Note : If $\lambda = \frac{1}{|\vec{a}|}$, provided that $\vec{a} \neq 0$, then $\lambda\vec{a}$ represents the unit vector in the direction of \vec{a} i.e. $\hat{a} = \frac{\vec{a}}{|\vec{a}|}$

COMPONENTS OF A VECTOR

▶ Let O be the origin and $P(x, y, z)$ be any point in space. Let $\hat{i}, \hat{j}, \hat{k}$ be unit vectors along the X -axis, Y -axis and Z -axis respectively. Then $\vec{OP} = x\hat{i} + y\hat{j} + z\hat{k}$, which is called the component form of \vec{OP} . Here x, y and z are scalar components of \vec{OP} and $x\hat{i}, y\hat{j}, z\hat{k}$ are vector components of \vec{OP} .

▶ If \vec{a} and \vec{b} are two given vectors as $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ and λ be any scalar, then

$$\vec{a} + \vec{b} = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}$$

$$\vec{a} - \vec{b} = (a_1 - b_1)\hat{i} + (a_2 - b_2)\hat{j} + (a_3 - b_3)\hat{k}$$

$$\lambda\vec{a} = (\lambda a_1)\hat{i} + (\lambda a_2)\hat{j} + (\lambda a_3)\hat{k}$$

$$\vec{a} = \vec{b} \Leftrightarrow a_1 = b_1, a_2 = b_2 \text{ and } a_3 = b_3$$

▶ \vec{a} and \vec{b} are collinear iff

$$\frac{b_1}{a_1} = \frac{b_2}{a_2} = \frac{b_3}{a_3} = \lambda.$$


VECTOR JOINING TWO POINTS

▶ If $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$ are any two points in the space then the vector joining P_1 and P_2 is the vector $\vec{P_1P_2}$.

Applying triangle law in ΔOP_1P_2 , we get

$$\vec{OP_1} + \vec{P_1P_2} = \vec{OP_2}$$

$$\Rightarrow \vec{P_1P_2} = \vec{OP_2} - \vec{OP_1}$$



$$= (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k})$$

$$= (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$$

$$\therefore |\vec{P_1P_2}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

SECTION FORMULA

▶ Let A, B be two points such that

$$\vec{OA} = \vec{a} \text{ and } \vec{OB} = \vec{b}.$$

▶ The position vector \vec{r} of the point P which divides the line segment AB internally in the

$$\text{ratio } m : n \text{ is given by } \vec{r} = \frac{m\vec{b} + n\vec{a}}{m+n}.$$

▶ The position vector \vec{r} of the point P which divides the line segment AB externally in the

$$\text{ratio } m : n \text{ is given by } \vec{r} = \frac{m\vec{b} - n\vec{a}}{m-n}.$$

▶ The position vector \vec{r} of the mid-point of the

$$\text{line segment } AB \text{ is given by } \vec{r} = \frac{\vec{a} + \vec{b}}{2}.$$

PRODUCT OF TWO VECTORS

▶ **Scalar (or dot) product :** The scalar (or dot) product of two (non-zero) vectors \vec{a} and \vec{b} , denoted by $\vec{a} \cdot \vec{b}$ (read as \vec{a} dot \vec{b}), is defined as $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = ab \cos \theta$,

where, $a = |\vec{a}|, b = |\vec{b}|$ and $\theta (0 \leq \theta \leq \pi)$ is the angle between \vec{a} and \vec{b} .

▶ **Properties of Scalar Product :**

(i) Scalar product is commutative: $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$

(ii) $\vec{a} \cdot \vec{0} = 0$

(iii) Scalar product is distributive over addition :

$$\bullet \vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$$

$$\bullet (\vec{a} + \vec{b}) \cdot \vec{c} = \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c}$$

(iv) $\lambda(\vec{a} \cdot \vec{b}) = (\lambda\vec{a}) \cdot \vec{b} = \vec{a} \cdot (\lambda\vec{b})$, λ be any scalar.

(v) If $\hat{i}, \hat{j}, \hat{k}$ are three unit vectors along three mutually perpendicular lines, then $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$ and $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$

(vi) Angle between two non-zero vectors

$$\vec{a} \text{ and } \vec{b} \text{ is given by } \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$


$$\text{i.e., } \theta = \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right)$$

(vii) Two non-zero vectors \vec{a} and \vec{b} are mutually perpendicular if and only if $\vec{a} \cdot \vec{b} = 0$

(viii) If $\theta = 0$, then $\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|$

If $\theta = \pi$, then $\vec{a} \cdot \vec{b} = -|\vec{a}||\vec{b}|$

►► **Projection of a vector on a line :** Let the vector \vec{AB} makes an angle θ with directed line ℓ .



Projection of \vec{AB} on $\ell = |\vec{AB}| \cos \theta = \vec{AC} = \vec{p}$.
The vector \vec{p} is called the projection vector. Its magnitude is $|\vec{p}|$, which is known as projection of vector \vec{AB} .

Projection of a vector \vec{a} on \vec{b} , is given as $\vec{a} \cdot \hat{b}$ i.e., $\frac{1}{|\vec{b}|}(\vec{a} \cdot \vec{b})$.

►► **Vector (or Cross) Product :** The vector (or cross) product of two (non-zero) vectors \vec{a} and \vec{b} (in an assigned order), denoted by $\vec{a} \times \vec{b}$ (read as \vec{a} cross \vec{b}), is defined as $\vec{a} \times \vec{b} = |\vec{a}||\vec{b}| \sin \theta \hat{n}$ where $\theta (0 \leq \theta \leq \pi)$ is the angle between \vec{a} and \vec{b} and \hat{n} is a unit vector perpendicular to both \vec{a} and \vec{b} .

► **Properties of Vector Product :**

(i) Non-commutative : $\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$

(ii) Vector product is distributive over addition :

$$\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$$

(iii) $\lambda(\vec{a} \times \vec{b}) = (\lambda\vec{a}) \times \vec{b} = \vec{a} \times (\lambda\vec{b})$, λ be any scalar.


(iv) $(\lambda_1\vec{a}) \times (\lambda_2\vec{b}) = \lambda_1\lambda_2(\vec{a} \times \vec{b})$

(v) $\hat{i} \times \hat{j} = \hat{k}$, $\hat{j} \times \hat{k} = \hat{i}$, $\hat{k} \times \hat{i} = \hat{j}$

(vi) Two non-zero vectors \vec{a} , \vec{b} are collinear if and only if $\vec{a} \times \vec{b} = \vec{0}$

Similarly, $\vec{a} \times \vec{a} = \vec{0}$ and $\vec{a} \times (-\vec{a}) = \vec{0}$, since in the first situation $\theta = 0$ and in the second one, $\theta = \pi$, making the value of $\sin \theta$ to be 0.


(vii) If \vec{a} and \vec{b} represent the adjacent sides of a triangle as given in the figure. Then,



$$\text{Area of triangle } ABC = \frac{1}{2} AB \cdot CD$$

$$= \frac{1}{2} |\vec{b}| |\vec{a}| \sin \theta = \frac{1}{2} |\vec{a} \times \vec{b}|$$

(viii) If \vec{a} and \vec{b} represent the adjacent sides of a parallelogram as given in the figure.



Then, area of parallelogram $ABCD = AB \cdot DE$

$$= |\vec{b}| |\vec{a}| \sin \theta = |\vec{a} \times \vec{b}|$$

(ix) If $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$, $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$= (a_2b_3 - a_3b_2)\hat{i} + (a_3b_1 - a_1b_3)\hat{j} + (a_1b_2 - a_2b_1)\hat{k}$$

(x) Angle between two vectors \vec{a} and \vec{b} is

$$\text{given by } \sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}||\vec{b}|}$$

$$\text{i.e., } \theta = \sin^{-1} \left(\frac{|\vec{a} \times \vec{b}|}{|\vec{a}||\vec{b}|} \right)$$

►► **Scalar Triple Product :** The scalar triple product of any three vectors \vec{a} , \vec{b} and \vec{c} is written as $\vec{a} \cdot (\vec{b} \times \vec{c})$ or $[\vec{a} \vec{b} \vec{c}]$.

► **Coplanarity of Three Vectors :** Three vectors \vec{a} , \vec{b} and \vec{c} are coplanar iff $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$.

► Volume of parallelepiped formed by adjacent sides given by the three vectors

$$\vec{a} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}), \quad \vec{b} = (b_1\hat{i} + b_2\hat{j} + b_3\hat{k}),$$

$$\text{and } \vec{c} = (c_1\hat{i} + c_2\hat{j} + c_3\hat{k}), \text{ is } |\vec{a} \cdot (\vec{b} \times \vec{c})|.$$

$$\text{i.e., } |\vec{a} \cdot (\vec{b} \times \vec{c})| = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

► For any three vectors \vec{a} , \vec{b} and \vec{c} ,

(i) $[\vec{a} \vec{b} \vec{c}] = [\vec{b} \vec{c} \vec{a}] = [\vec{c} \vec{a} \vec{b}]$

(ii) $[\vec{a} \vec{b} \vec{c}] = -[\vec{a} \vec{c} \vec{b}]$ (iii) $[\vec{a} \vec{a} \vec{b}] = 0$

Previous Years' CBSE Board Questions

10.2 Some Basic Concepts

VSA (1 mark)

- Find a vector \vec{a} of magnitude $5\sqrt{2}$, making an angle of $\frac{\pi}{4}$ with x -axis, $\frac{\pi}{2}$ with y -axis and an acute angle θ with z -axis. (AI 2014)
- If a unit vector \vec{a} makes angles $\frac{\pi}{3}$ with \hat{i} , $\frac{\pi}{4}$ with \hat{j} and an acute angle θ with \hat{k} , then find the value of θ . (Delhi 2013)
- Find the magnitude of the vector $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k}$. (AI 2011C)

10.3 Types of Vectors

VSA (1 mark)

- The value of p for which $p(\hat{i} + \hat{j} + \hat{k})$ is a unit vector is
(a) 0 (b) $\frac{1}{\sqrt{3}}$
(c) 1 (d) $\sqrt{3}$ (2020)

10.4 Addition of Vectors

VSA (1 mark)

- $ABCD$ is a rhombus, whose diagonals intersect at E . Then $\vec{EA} + \vec{EB} + \vec{EC} + \vec{ED}$ equals
(a) $\vec{0}$ (b) \vec{AD}
(c) $2\vec{BC}$ (d) $2\vec{AD}$ (2020)
- Find the sum of the vectors $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$ and $\vec{c} = \hat{i} - 6\hat{j} - 7\hat{k}$. (Delhi 2012)
- Find the sum of the following vectors :
 $\vec{a} = \hat{i} - 3\hat{k}$, $\vec{b} = 2\hat{j} - \hat{k}$, $\vec{c} = 2\hat{i} - 3\hat{j} + 2\hat{k}$
(Delhi 2012)
- Find the sum of the following vectors :
 $\vec{a} = \hat{i} - 2\hat{j}$, $\vec{b} = 2\hat{i} - 3\hat{j}$, $\vec{c} = 2\hat{i} + 3\hat{k}$ (Delhi 2012)

- If A, B and C are the vertices of a triangle ABC , then what is the value of $\vec{AB} + \vec{BC} + \vec{CA}$?

(Delhi 2011C)

10.5 Multiplication of a Vector by a Scalar

VSA (1 mark)

- The position vector of two points A and B are $\vec{OA} = 2\hat{i} - \hat{j} - \hat{k}$ and $\vec{OB} = 2\hat{i} - \hat{j} + 2\hat{k}$, respectively. The position vector of a point P which divides the line segment joining A and B in the ratio $2 : 1$ is _____. (2020)
- Find the position vector of a point which divides the join of points with position vectors $\vec{a} - 2\vec{b}$ and $2\vec{a} + \vec{b}$ externally in the ratio $2 : 1$. (Delhi 2016)
- Write the position vector of the point which divides the join of points with position vectors $3\vec{a} - 2\vec{b}$ and $2\vec{a} + 3\vec{b}$ in the ratio $2 : 1$. (AI 2016)
- Find the unit vector in the direction of the sum of the vectors $2\hat{i} + 3\hat{j} - \hat{k}$ and $4\hat{i} - 3\hat{j} + 2\hat{k}$. (Foreign 2015)
- Find a vector in the direction of $\vec{a} = \hat{i} - 2\hat{j}$ that has magnitude 7 units. (Delhi 2015C)
- Write the direction ratios of the vector $3\vec{a} + 2\vec{b}$ where $\vec{a} = \hat{i} + \hat{j} - 2\hat{k}$ and $\vec{b} = 2\hat{i} - 4\hat{j} + 5\hat{k}$. (AI 2015C)
- Write a unit vector in the direction of the sum of the vectors $\vec{a} = 2\hat{i} + 2\hat{j} - 5\hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} - 7\hat{k}$. (Delhi 2014)
- Find the value of ' p ' for which the vectors $3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\hat{i} - 2p\hat{j} + 3\hat{k}$ are parallel. (AI 2014)
- Find a vector in the direction of vector $2\hat{i} - 3\hat{j} + 6\hat{k}$ which has magnitude 21 units. (Foreign 2014)

19. Write a unit vector in the direction of vector \overline{PQ} , where P and Q are the points $(1, 3, 0)$ and $(4, 5, 6)$ respectively. (Foreign 2014)
20. Write a vector in the direction of the vector $\hat{i} - 2\hat{j} + 2\hat{k}$ that has magnitude 9 units. (Delhi 2014C)
21. If $\vec{a} = x\hat{i} + 2\hat{j} - z\hat{k}$ and $\vec{b} = 3\hat{i} - y\hat{j} + \hat{k}$ are two equal vectors, then write the value of $x + y + z$. (Delhi 2013)
22. Write a unit vector in the direction of the sum of vectors $\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + \hat{j} + 3\hat{k}$. (Delhi 2013)
23. P and Q are two points with position vectors $3\vec{a} - 2\vec{b}$ and $\vec{a} + \vec{b}$ respectively. Write the position vector of a point R which divides the line segment PQ in the ratio $2 : 1$ externally. (AI 2013)
24. A and B are two points with position vectors $2\vec{a} - 3\vec{b}$ and $6\vec{b} - \vec{a}$ respectively. Write the position vector of a point P which divides the line segment AB internally in the ratio $1 : 2$. (AI 2013)
25. L and M are two points with position vectors $2\vec{a} - \vec{b}$ and $\vec{a} + 2\vec{b}$ respectively. Write the position vector of a point N which divides the line segment LM in the ratio $2 : 1$ externally. (AI 2013)
26. Find the scalar components of the vector \overline{AB} with initial point $A(2, 1)$ and terminal point $B(-5, 7)$. (AI 2012)
27. Find a unit vector parallel to the sum of the vectors $\hat{i} + \hat{j} + \hat{k}$ and $2\hat{i} - 3\hat{j} + 5\hat{k}$. (Delhi 2012C)
28. Find a unit vector in the direction of the vector $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k}$. (AI 2012C)
29. Write the direction cosines of the vector $-2\hat{i} + \hat{j} - 5\hat{k}$. (Delhi 2011)
30. For what value of 'a', the vectors $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $a\hat{i} + 6\hat{j} - 8\hat{k}$ are collinear? (Delhi 2011)
31. Write a unit vector in the direction of the vector $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$. (AI 2011)
32. Find a unit vector in the direction of $\vec{a} = 2\hat{i} - 3\hat{j} + 6\hat{k}$. (Delhi 2011C)
33. Find a unit vector in the direction of the vector $\vec{a} = 2\hat{i} + 3\hat{j} + 6\hat{k}$. (AI 2011C)

SA (2 marks)

34. X and Y are two points with position vectors $3\vec{a} + \vec{b}$ and $\vec{a} - 3\vec{b}$ respectively. Write the position vector of a point Z which divides the line segment XY in the ratio $2 : 1$ externally. (AI 2019)

LA 1 (4 marks)

35. The two vectors $\hat{j} + \hat{k}$ and $3\hat{i} - \hat{j} + 4\hat{k}$ represent the two sides AB and AC , respectively of a ΔABC . Find the length of the median through A . (Delhi 2016, Foreign 2015)
36. Find a vector of magnitude 5 units and parallel to the resultant of the vectors $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$. (Delhi 2011)

10.6 Product of Two Vectors

VSA (1 mark)

37. If the projection of $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$ on $\vec{b} = 2\hat{i} + \lambda\hat{k}$ is zero, then the value of λ is
(a) 0 (b) 1
(c) $\frac{-2}{3}$ (d) $\frac{-3}{2}$ (2020)
38. The area of the parallelogram whose diagonals are $2\hat{i}$ and $-3\hat{k}$ is _____ square units.
39. The value of λ for which the vectors $2\hat{i} - \lambda\hat{j} + \hat{k}$ and $\hat{i} + 2\hat{j} - \hat{k}$ are orthogonal is _____. (2020)
40. If $\hat{i}, \hat{j}, \hat{k}$ are unit vectors along three mutually perpendicular directions, then
(a) $\hat{i} \cdot \hat{j} = 1$ (b) $\hat{i} \times \hat{j} = 1$
(c) $\hat{i} \cdot \hat{k} = 0$ (d) $\hat{i} \times \hat{k} = 0$ (2020)

41. Find the magnitude of each of the two vectors \vec{a} and \vec{b} , having the same magnitude such that the angle between them is 60° and their scalar product is $\frac{9}{2}$. (2018)
42. Write the number of vectors of unit length perpendicular to both the vectors $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = \hat{j} + \hat{k}$. (AI 2016)
43. If $\vec{a}, \vec{b}, \vec{c}$ are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, then write the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$. (Foreign 2016)
44. If $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 400$ and $|\vec{a}| = 5$ then write the value of $|\vec{b}|$. (Foreign 2016)
45. If $\vec{a} = 7\hat{i} + \hat{j} - 4\hat{k}$ and $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$, then find the projection of \vec{a} on \vec{b} . (Delhi 2015, 2013C)
46. If \hat{a}, \hat{b} and \hat{c} are mutually perpendicular unit vectors, then find the value of $|2\hat{a} + \hat{b} + \hat{c}|$. (AI 2015)
47. Write a unit vector perpendicular to both the vectors $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$. (AI 2015)
48. Find the area of a parallelogram whose adjacent sides are represented by the vectors $2\hat{i} - 3\hat{k}$ and $4\hat{j} + 2\hat{k}$. (Foreign 2015)
49. If \vec{a} and \vec{b} are unit vectors, then what is the angle between \vec{a} and \vec{b} so that $\sqrt{2}\vec{a} - \vec{b}$ is a unit vector? (Delhi 2015C)
50. Find the projection of the vector $\vec{a} = 2\hat{i} + 3\hat{j} + 2\hat{k}$ on the vector $\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$. (AI 2015C)
51. Find the projection of vector $\hat{i} + 3\hat{j} + 7\hat{k}$ on the vector $2\hat{i} - 3\hat{j} + 6\hat{k}$. (Delhi 2014)
52. If \vec{a} and \vec{b} are two unit vectors such that $\vec{a} + \vec{b}$ is also a unit vector, then find the angle between \vec{a} and \vec{b} . (Delhi 2014)
53. If vectors \vec{a} and \vec{b} are such that, $|\vec{a}| = 3, |\vec{b}| = \frac{2}{3}$ and $\vec{a} \times \vec{b}$ is a unit vector, then write the angle between \vec{a} and \vec{b} . (Delhi 2014)
54. If \vec{a} and \vec{b} are perpendicular vectors, $|\vec{a} + \vec{b}| = 13$ and $|\vec{a}| = 5$, find the value of $|\vec{b}|$. (AI 2014)
55. Write the projection of the vector $\hat{i} + \hat{j} + \hat{k}$ along the vector \hat{j} . (Foreign 2014)
56. Write the value of $\hat{i} \times (\hat{j} + \hat{k}) + \hat{j} \times (\hat{k} + \hat{i}) + \hat{k} \times (\hat{i} + \hat{j})$. (Foreign 2014)
57. Write the projection of the vector $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ on the vector $\vec{b} = \hat{i} + 2\hat{j} + 2\hat{k}$. (Delhi 2014C)
58. If \vec{a} and \vec{b} are unit vectors, then find the angle between \vec{a} and \vec{b} , given that $(\sqrt{3}\vec{a} - \vec{b})$ is a unit vector. (Delhi 2014C)
59. Write the value of cosine of the angle which the vector $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ makes with y -axis. (Delhi 2014C)
60. If $|\vec{a}| = 8, |\vec{b}| = 3$ and $|\vec{a} \times \vec{b}| = 12$, find the angle between \vec{a} and \vec{b} . (AI 2014C)
1. Find the angle between x -axis and the vector $\hat{i} + \hat{j} + \hat{k}$. (AI 2014C)
62. Find $|\vec{x}|$, if for a unit vector \vec{a} , $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 15$ (AI 2013)
63. Write the value of λ so that the vectors $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + 3\hat{k}$ are perpendicular to each other. (Delhi 2013C, AI 2012C)
64. For what value of λ are the vectors $\hat{i} + 2\lambda\hat{j} + \hat{k}$ and $2\hat{i} + \hat{j} - 3\hat{k}$ are perpendicular? (AI 2013C, 2011C, Delhi 2012C)
65. Write the projection of $\vec{b} + \vec{c}$ on \vec{a} , where $\vec{a} = 2\hat{i} - 2\hat{j} + \hat{k}, \vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$. (AI 2013C)
66. Find ' λ ' when the projection of $\vec{a} = \lambda\hat{i} + \hat{j} + 4\hat{k}$ on $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$ is 4 units. (Delhi 2012)
67. Write the value of $(\hat{i} \times \hat{j}) \cdot \hat{k} + \hat{i} \cdot \hat{j}$. (AI 2012)

68. Write the value of $(\hat{k} \times \hat{j}) \cdot \hat{i} + \hat{j} \cdot \hat{k}$. (AI 2012)
69. Write the value of $(\hat{k} \times \hat{i}) \cdot \hat{j} + \hat{i} \cdot \hat{k}$. (AI 2012)
70. Write the angle between two vectors \vec{a} and \vec{b} with magnitudes $\sqrt{3}$ and 2 respectively having $\vec{a} \cdot \vec{b} = \sqrt{6}$. (AI 2011)
71. Write the projection of the vector $\hat{i} - \hat{j}$ on the vector $\hat{i} + \hat{j}$. (AI 2011)
72. If $|\vec{a}| = \sqrt{3}, |\vec{b}| = 2$ and angle between \vec{a} and \vec{b} is 60° , find $\vec{a} \cdot \vec{b}$. (Delhi 2011C)

SA (2 marks)

73. Find a unit vector perpendicular to each of the vectors \vec{a} and \vec{b} and where $\vec{a} = 5\hat{i} + 6\hat{j} - 2\hat{k}$ and $\vec{b} = 7\hat{i} + 6\hat{j} + 2\hat{k}$. (2020)
74. Show that for any two non-zero vectors \vec{a} and \vec{b} , $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ iff \vec{a} and \vec{b} are perpendicular vectors. (2020)
75. Show that the vectors $2\hat{i} - \hat{j} + \hat{k}$, $3\hat{i} + 7\hat{j} + \hat{k}$ and $5\hat{i} + 6\hat{j} + 2\hat{k}$ form the sides of a right-angled triangle. (2020)
76. If the sum of two unit vectors is a unit vector, prove that the magnitude of their difference is $\sqrt{3}$. (Delhi 2019)
77. Let $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$ and $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$ be two vectors. Show that the vectors $(\vec{a} + \vec{b})$ and $(\vec{a} - \vec{b})$ are perpendicular to each other. (AI 2019)
78. If θ is the angle between two vectors $\hat{i} - 2\hat{j} + 3\hat{k}$ and $3\hat{i} - 2\hat{j} + \hat{k}$, find $\sin \theta$. (2018)

LA 1 (4 marks)

79. If $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$ and $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$ represent two adjacent sides of a parallelogram, find unit vectors parallel to the diagonals of the parallelogram. (2020)
80. Using vectors, find the area of the triangle ABC with vertices $A(1, 2, 3)$, $B(2, -1, 4)$ and $C(4, 5, -1)$ (2020, Delhi 2013, AI 2013)
81. If $\hat{i} + \hat{j} + \hat{k}$, $2\hat{i} + 5\hat{j}$, $3\hat{i} + 2\hat{j} - 3\hat{k}$ and $\hat{i} - 6\hat{j} - \hat{k}$ respectively are the position vectors of points A, B, C and D, then find the angle between the straight lines AB and CD. Find whether \overline{AB} and \overline{CD} are collinear or not. (Delhi 2019)
82. Let $\vec{a} = 4\hat{i} + 5\hat{j} - \hat{k}$, $\vec{b} = \hat{i} - 4\hat{j} + 5\hat{k}$ and $\vec{c} = 3\hat{i} + \hat{j} - \hat{k}$. Find a vector \vec{d} which is perpendicular to both \vec{c} and \vec{b} and $\vec{d} \cdot \vec{a} = 21$. (2018)
83. If \vec{a} , \vec{b} , \vec{c} are mutually perpendicular vectors of equal magnitudes, show that the vector $\vec{a} + \vec{b} + \vec{c}$ is equally inclined to \vec{a} , \vec{b} and \vec{c} . Also, find the angle which $\vec{a} + \vec{b} + \vec{c}$ makes with \vec{a} or \vec{b} or \vec{c} . (Delhi 2017)
84. Show that the points A, B, C with position vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $3\hat{i} - 4\hat{j} - 4\hat{k}$ respectively, are the vertices of a right-angled triangle. Hence find the area of the triangle. (AI 2017)
85. The two adjacent sides of a parallelogram are $2\hat{i} - 4\hat{j} - 5\hat{k}$ and $2\hat{i} + 2\hat{j} + 3\hat{k}$. Find the two unit vectors parallel to its diagonals. Using the diagonal vectors, find the area of the parallelogram. (AI 2016)
86. If $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$ and $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$, show that $\vec{a} - \vec{d}$ is parallel to $\vec{b} - \vec{c}$, where $\vec{a} \neq \vec{d}$ and $\vec{b} \neq \vec{c}$. (Foreign 2016)
87. If $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, find $(\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j}) + xy$. (Delhi 2015)
88. If $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j}$ and $\vec{c} = 3\hat{i} - 4\hat{j} - 5\hat{k}$, then find a unit vector perpendicular to both of the vectors $(\vec{a} - \vec{b})$ and $(\vec{c} - \vec{b})$. (AI 2015)
89. Vectors \vec{a} , \vec{b} and \vec{c} are such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ and $|\vec{a}| = 3$, $|\vec{b}| = 5$ and $|\vec{c}| = 7$. Find the angle between \vec{a} and \vec{b} . (Delhi 2014)
90. The scalar product of the vector $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ with a unit vector along the sum of vectors $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$ and $\vec{c} = \lambda\hat{i} + 2\hat{j} + 3\hat{k}$ is

- equal to one. Find the value of λ and hence find the unit vector along $\vec{b} + \vec{c}$. (AI 2014)
91. Find a unit vector perpendicular to both of the vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ where $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$. (Foreign 2014)
92. If $\vec{a} = 2\hat{i} - 3\hat{j} + \hat{k}, \vec{b} = -\hat{i} + \hat{k}, \vec{c} = 2\hat{j} - \hat{k}$ are three vectors, find the area of the parallelogram having diagonals $(\vec{a} + \vec{b})$ and $(\vec{b} + \vec{c})$. (Delhi 2014C)
93. Find the vector \vec{p} which is perpendicular to both $\vec{\alpha} = 4\hat{i} + 5\hat{j} - \hat{k}$ and $\vec{\beta} = \hat{i} - 4\hat{j} + 5\hat{k}$ and $\vec{p} \cdot \vec{q} = 21$, where $\vec{q} = 3\hat{i} + \hat{j} - \hat{k}$. (AI 2014C)
94. If \vec{a} and \vec{b} are two vectors such that $|\vec{a} + \vec{b}| = |\vec{a}|$, then prove that vector $2\vec{a} + \vec{b}$ is perpendicular to vector \vec{b} . (Delhi 2013)
95. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = \hat{j} - \hat{k}$, find a vector \vec{c} , such that $\vec{a} \times \vec{c} = \vec{b}$ and $\vec{a} \cdot \vec{c} = 3$. (Delhi 2013)
96. If $\vec{a} = \hat{i} - \hat{j} + 7\hat{k}$ and $\vec{b} = 5\hat{i} - \hat{j} + \lambda\hat{k}$, then find the value of λ , so that $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ are perpendicular vectors. (AI 2013)
97. If $\vec{a}, \vec{b}, \vec{c}$ are three mutually perpendicular vectors of the same magnitude, prove that $\vec{a} + \vec{b} + \vec{c}$ is equally inclined with the vectors \vec{a}, \vec{b} and \vec{c} . (Delhi 2013C)
98. Dot product of a vector with vectors $\hat{i} - \hat{j} + \hat{k}, 2\hat{i} + \hat{j} - 3\hat{k}$ and $\hat{i} + \hat{j} + \hat{k}$ are respectively 4, 0 and 2. Find the vector. (Delhi 2013C)
99. Find the values of λ for which the angle between the vectors $\vec{a} = 2\lambda^2 \hat{i} + 4\lambda\hat{j} + \hat{k}$ and $\vec{b} = 7\hat{i} - 2\hat{j} + \lambda\hat{k}$ is obtuse. (AI 2013C)
100. If \vec{a}, \vec{b} and \vec{c} are three vectors such that each one is perpendicular to the vector obtained by sum of the other two and $|\vec{a}| = 3, |\vec{b}| = 4$ and $|\vec{c}| = 5$, then prove that $|\vec{a} + \vec{b} + \vec{c}| = 5\sqrt{2}$. (AI 2013C)
101. If $\vec{a} = 3\hat{i} - \hat{j}$ and $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$ then express \vec{b} in the form $\vec{b} = \vec{b}_1 + \vec{b}_2$ where $\vec{b}_1 \parallel \vec{a}$ and $\vec{b}_2 \perp \vec{a}$. (AI 2013C)
102. If $\vec{a}, \vec{b}, \vec{c}$ are three vectors such that $|\vec{a}| = 5, |\vec{b}| = 12$ and $|\vec{c}| = 13$ and $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, find the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$. (Delhi 2012)
103. Let $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$. Find a vector \vec{p} which is perpendicular to both \vec{a} and \vec{b} and $\vec{p} \cdot \vec{c} = 18$. (AI 2012)
104. If the sum of two unit vectors \hat{a} and \hat{b} is a unit vector, show that the magnitude of their difference is $\sqrt{3}$. (Delhi 2012C)
105. Find a unit vector perpendicular to each of the vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$, where $\vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$. (Delhi 2011)
106. If two vectors \vec{a} and \vec{b} are such that $|\vec{a}| = 2, |\vec{b}| = 1$ and $\vec{a} \cdot \vec{b} = 1$, then find the value of $(3\vec{a} - 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$. (Delhi 2011)
107. Using vectors, find the area of the triangle with vertices A(1, 1, 2), B(2, 3, 5) and C(1, 5, 5). (AI 2011)
108. If \vec{a}, \vec{b} and \vec{c} are three mutually perpendicular vectors of equal magnitude, show that $\vec{a} + \vec{b} + \vec{c}$ is equally inclined to \vec{a}, \vec{b} and \vec{c} . Also find the angle. (Delhi 2011C)
109. If \vec{a}, \vec{b} and \vec{c} are three vectors such that $|\vec{a}| = 3, |\vec{b}| = 4$ and $|\vec{c}| = 5$ and each one of them is perpendicular to the sum of the other two, then find $|\vec{a} + \vec{b} + \vec{c}|$. (AI 2011C)

10.7 Scalar Triple Product

VSA (1 mark)

110. Find λ , if the vectors $\vec{a} = \hat{i} + 3\hat{j} + \hat{k}, \vec{b} = 2\hat{i} - \hat{j} - \hat{k}$ and $\vec{c} = \lambda\hat{j} + 3\hat{k}$ are coplanar. (Delhi 2015)
111. Find $\vec{a} \cdot (\vec{b} \times \vec{c})$, if $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 3\hat{i} + \hat{j} + 2\hat{k}$. (AI 2014)

SA (2 marks)

112. Find the volume of the parallelepiped whose adjacent edges are represented by $2\vec{a}$, $-\vec{b}$ and $3\vec{c}$, where
 $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} + 4\hat{j} - 5\hat{k}$ and
 $\vec{c} = 2\hat{i} - \hat{j} + 3\hat{k}$. (2020)
113. If $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$ and
 $\vec{c} = -3\hat{i} + \hat{j} + 2\hat{k}$, find $[\vec{a} \vec{b} \vec{c}]$. (Delhi 2019)

LA 1 (4 marks)

114. Find the value of x , for which the four points $A(x, -1, -1)$, $B(4, 5, 1)$, $C(3, 9, 4)$ and $D(-4, 4, 4)$ are coplanar. (AI 2019)
115. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i}$ and $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$, then
 (a) Let $c_1 = 1$ and $c_2 = 2$, find c_3 which makes \vec{a} , \vec{b} and \vec{c} coplanar.
 (b) If $c_2 = -1$ and $c_3 = 1$, show that no value of c_1 can make \vec{a} , \vec{b} and \vec{c} coplanar. (Delhi 2017)
116. Find the value of λ , if four points with position vectors $3\hat{i} + 6\hat{j} + 9\hat{k}$, $\hat{i} + 2\hat{j} + 3\hat{k}$, $2\hat{i} + 3\hat{j} + \hat{k}$ and $4\hat{i} + 6\hat{j} + \lambda\hat{k}$ are coplanar. (AI 2017)

117. Show that the vectors \vec{a} , \vec{b} and \vec{c} are coplanar if $\vec{a} + \vec{b}$, $\vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$ are coplanar. (Delhi 2016)
118. Find the value of λ so that the four points A , B , C and D with position vectors $4\hat{i} + 5\hat{j} + \hat{k}$, $-\hat{j} - \hat{k}$, $3\hat{i} + \lambda\hat{j} + 4\hat{k}$ and $-4\hat{i} + 4\hat{j} + 4\hat{k}$ respectively are coplanar. (Delhi 2015C)
119. Prove that : $[\vec{a}, \vec{b} + \vec{c}, \vec{d}] = [\vec{a}, \vec{b}, \vec{d}] + [\vec{a}, \vec{c}, \vec{d}]$. (AI 2015C)
120. Prove that, for any three vectors \vec{a} , \vec{b} , \vec{c}
 $[\vec{a} + \vec{b} \vec{b} + \vec{c} \vec{c} + \vec{a}] = 2[\vec{a} \vec{b} \vec{c}]$ (Delhi 2014)
121. Show that the four points A , B , C and D with position vectors $4\hat{i} + 5\hat{j} + \hat{k}$, $-\hat{j} - \hat{k}$, $3\hat{i} + 9\hat{j} + 4\hat{k}$ and $4(-\hat{i} + \hat{j} + \hat{k})$ respectively are coplanar. (AI 2014)
122. Show that the vectors $\vec{a}, \vec{b}, \vec{c}$ are coplanar if and only if $\vec{a} + \vec{b}, \vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$ are coplanar. (Foreign 2014)
123. If the three vectors \vec{a}, \vec{b} and \vec{c} are coplanar, prove that the vectors $\vec{a} + \vec{b}, \vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$ are also coplanar. (Delhi 2014C, 2013C)

Detailed Solutions

1. Here, $l = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$, $m = \cos \frac{\pi}{2} = 0$,
 $n = \cos \theta$
 Since, $l^2 + m^2 + n^2 = 1$
 $\Rightarrow \frac{1}{2} + 0 + \cos^2 \theta = 1 \Rightarrow \cos^2 \theta = 1 - \frac{1}{2} = \frac{1}{2}$
 $\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4} \therefore n = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$
 \therefore The vector of magnitude $5\sqrt{2}$ is
 $\vec{a} = 5\sqrt{2}(\hat{i} + \hat{j} + n\hat{k})$
 $= 5\sqrt{2}\left(\frac{1}{\sqrt{2}}\hat{i} + 0\hat{j} + \frac{1}{\sqrt{2}}\hat{k}\right) = 5(\hat{i} + \hat{k})$
2. $l = \cos \frac{\pi}{3} = \frac{1}{2}$, $m = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$ and $n = \cos \theta$

- Now, $l^2 + m^2 + n^2 = 1$
 $\Rightarrow \left(\frac{1}{2}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2 + n^2 = 1$
 $\Rightarrow \frac{1}{4} + \frac{1}{2} + n^2 = 1 \Rightarrow n^2 = \frac{1}{4} \Rightarrow n = \pm \frac{1}{2}$
 $\Rightarrow \cos \theta = \pm \frac{1}{2}$
 But θ is an acute angle (given).
 $\therefore \theta = \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3}$
3. Here, $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k}$
 \therefore Its magnitude = $|\vec{a}|$
 $= \sqrt{3^2 + (-2)^2 + 6^2} = \sqrt{9 + 4 + 36} = \sqrt{49} = 7$.

4. (b) : Let $a = (\hat{i} + \hat{j} + \hat{k})$

So, unit vector of $\vec{a} = \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{1+1+1}} = \frac{1}{\sqrt{3}}(\hat{i} + \hat{j} + \hat{k})$

\therefore The value of p is $\frac{1}{\sqrt{3}}$.

5. (a) : $\vec{EA} + \vec{EB} + \vec{EC} + \vec{ED}$
 $= \vec{EA} + \vec{EB} - \vec{EA} - \vec{EB}$

[As diagonals of a rhombus bisect each other]
 $= \vec{0}$

6. The given vectors are

$\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$, $\vec{c} = \hat{i} - 6\hat{j} - 7\hat{k}$

\therefore Their sum $= \vec{a} + \vec{b} + \vec{c}$
 $= (\hat{i} - 2\hat{j} + \hat{k}) + (-2\hat{i} + 4\hat{j} + 5\hat{k}) + (\hat{i} - 6\hat{j} - 7\hat{k})$
 $= -4\hat{j} - \hat{k}$.


7. Required sum $= \vec{a} + \vec{b} + \vec{c}$
 $= (\hat{i} - 3\hat{k}) + (2\hat{j} - \hat{k}) + (2\hat{i} - 3\hat{j} + 2\hat{k})$
 $= 3\hat{i} - \hat{j} - 2\hat{k}$.

8. Required sum $= \vec{a} + \vec{b} + \vec{c}$
 $= (\hat{i} - 2\hat{j}) + (2\hat{i} - 3\hat{j}) + (2\hat{i} + 3\hat{k}) = 5\hat{i} - 5\hat{j} + 3\hat{k}$.

9. Let ABC be the given triangle.

Now $\vec{AB} + \vec{BC} = \vec{AC}$
(By Triangle law)

$\Rightarrow \vec{AB} + \vec{BC} + \vec{CA} = \vec{AC} + \vec{CA} = \vec{0}$



10. Required position vector of point P

$$= \frac{1(2\hat{i} - \hat{j} - \hat{k}) + 2(2\hat{i} - \hat{j} + 2\hat{k})}{2+1}$$

$$= \frac{2\hat{i} - \hat{j} - \hat{k} + 4\hat{i} - 2\hat{j} + 4\hat{k}}{3}$$

$$= \frac{1}{3}(6\hat{i} - 3\hat{j} + 3\hat{k}) = 2\hat{i} - \hat{j} + \hat{k}$$

11. Required position vector

$$= \frac{2 \cdot (2\vec{a} + \vec{b}) - 1(\vec{a} - 2\vec{b})}{2-1} = \frac{4\vec{a} + 2\vec{b} - \vec{a} + 2\vec{b}}{1}$$

$$= 3\vec{a} + 4\vec{b}$$

12. Required position vector

$$= \frac{2(2\vec{a} + 3\vec{b}) + 1(3\vec{a} - 2\vec{b})}{2+1} = \frac{7\vec{a} + 4\vec{b}}{3} = \frac{7}{3}\vec{a} + \frac{4}{3}\vec{b}$$

13. Let $a = 2\hat{i} + 3\hat{j} - \hat{k}$ and $b = 4\hat{i} - 3\hat{j} + 2\hat{k}$.

Then, the sum of the given vectors is

$\vec{c} = \vec{a} + \vec{b} = (2+4)\hat{i} + (3-3)\hat{j} + (-1+2)\hat{k} = 6\hat{i} + \hat{k}$

and $|\vec{c}| = |\vec{a} + \vec{b}| = \sqrt{6^2 + 1^2} = \sqrt{36+1} = \sqrt{37}$

\therefore Unit vector, $\hat{c} = \frac{\vec{c}}{|\vec{c}|} = \frac{6\hat{i} + \hat{k}}{\sqrt{37}} = \frac{6}{\sqrt{37}}\hat{i} + \frac{1}{\sqrt{37}}\hat{k}$

14. A unit vector in the direction of $\vec{a} = \hat{i} - 2\hat{j}$

is $\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{\hat{i} - 2\hat{j}}{\sqrt{1^2 + (-2)^2}} = \frac{1}{\sqrt{5}}(\hat{i} - 2\hat{j})$

\therefore The required vector of magnitude 7 in the direction of $\vec{a} = 7 \cdot \hat{a} = \frac{7}{\sqrt{5}}(\hat{i} - 2\hat{j})$.

15. $\vec{a} = \hat{i} + \hat{j} - 2\hat{k}$; $\vec{b} = 2\hat{i} - 4\hat{j} + 5\hat{k}$

$\therefore 3\vec{a} + 2\vec{b} = 3(\hat{i} + \hat{j} - 2\hat{k}) + 2(2\hat{i} - 4\hat{j} + 5\hat{k})$
 $= (3\hat{i} + 3\hat{j} - 6\hat{k}) + (4\hat{i} - 8\hat{j} + 10\hat{k}) = 7\hat{i} - 5\hat{j} + 4\hat{k}$

\therefore The direction ratios of the vector $3\vec{a} + 2\vec{b}$ are 7, -5, 4.

16. Refer to answer 13.

17. Let $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\vec{b} = \hat{i} - 2p\hat{j} + 3\hat{k}$

For \vec{a} and \vec{b} to be parallel, $\vec{b} = \lambda\vec{a}$.

$\Rightarrow \hat{i} - 2p\hat{j} + 3\hat{k} = \lambda(3\hat{i} + 2\hat{j} + 9\hat{k}) = 3\lambda\hat{i} + 2\lambda\hat{j} + 9\lambda\hat{k}$

$\Rightarrow 1 = 3\lambda$; $-2p = 2\lambda$, $3 = 9\lambda$

$\Rightarrow \lambda = \frac{1}{3}$ and $p = -\lambda = -\frac{1}{3}$

18. Let $\vec{a} = 2\hat{i} - 3\hat{j} + 6\hat{k}$.

The vector in the direction of \vec{a} with a magnitude of 21 $= 21 \times \hat{a}$

\therefore Required vector $= 21 \times \frac{2\hat{i} - 3\hat{j} + 6\hat{k}}{\sqrt{2^2 + (-3)^2 + 6^2}}$

$= 21 \times \frac{2\hat{i} - 3\hat{j} + 6\hat{k}}{7} = 6\hat{i} - 9\hat{j} + 18\hat{k}$

19. We have $\vec{PQ} = \vec{OQ} - \vec{OP}$

$= (4\hat{i} + 5\hat{j} + 6\hat{k}) - (\hat{i} + 3\hat{j}) = 3\hat{i} + 2\hat{j} + 6\hat{k}$

Required unit vector $= \frac{3\hat{i} + 2\hat{j} + 6\hat{k}}{\sqrt{3^2 + 2^2 + 6^2}} = \frac{3\hat{i} + 2\hat{j} + 6\hat{k}}{7}$

20. Refer to answer 18.

21. Given, $\vec{a} = \vec{b}$

$$\Rightarrow x\hat{i} + 2\hat{j} - z\hat{k} = 3\hat{i} - y\hat{j} + \hat{k}$$

$$\therefore x = 3, y = -2, z = -1$$

Hence, the value of $x + y + z = 0$

22. Refer to answer 13.

23. Refer to answer 11.

24. Refer to answer 12.

25. Refer to answer 11.

26. Vector $\vec{AB} = \vec{OB} - \vec{OA}$

$$= (-5\hat{i} + 7\hat{j}) - (2\hat{i} + \hat{j}) = -7\hat{i} + 6\hat{j}$$

So, its scalar components are $(-7, 6)$.

27. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} - 3\hat{j} + 5\hat{k}$

$$\therefore \vec{a} + \vec{b} = (\hat{i} + \hat{j} + \hat{k}) + (2\hat{i} - 3\hat{j} + 5\hat{k})$$

$$= 3\hat{i} - 2\hat{j} + 6\hat{k}$$

Any vector parallel to $\vec{a} + \vec{b}$

$$= \lambda(\vec{a} + \vec{b}) = \lambda(3\hat{i} - 2\hat{j} + 6\hat{k})$$

\therefore The unit vector in this direction

$$= \frac{\lambda(3\hat{i} - 2\hat{j} + 6\hat{k})}{\sqrt{(3\lambda)^2 + (-2\lambda)^2 + (6\lambda)^2}}$$

$$= \frac{\lambda(3\hat{i} - 2\hat{j} + 6\hat{k})}{|\lambda| \cdot 7} = \pm \frac{1}{7}(3\hat{i} - 2\hat{j} + 6\hat{k})$$

28. The given vector is $\vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k}$

$$\Rightarrow |\vec{a}| = \sqrt{3^2 + (-2)^2 + 6^2} = 7$$

\therefore A unit vector in the direction of vector \vec{a} is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{1}{7}(3\hat{i} - 2\hat{j} + 6\hat{k})$$

29. We have, $\vec{a} = -2\hat{i} + \hat{j} - 5\hat{k}$

Direction cosines of the given vector are

$$\left(\frac{-2}{\sqrt{(-2)^2 + (1)^2 + (-5)^2}}, \frac{1}{\sqrt{(-2)^2 + (1)^2 + (-5)^2}}, \frac{-5}{\sqrt{(-2)^2 + (1)^2 + (-5)^2}} \right)$$

$$= \left(\frac{-2}{\sqrt{4+1+25}}, \frac{1}{\sqrt{4+1+25}}, \frac{-5}{\sqrt{4+1+25}} \right)$$

$$\therefore \text{Direction cosines are } \left(\frac{-2}{\sqrt{30}}, \frac{1}{\sqrt{30}}, \frac{-5}{\sqrt{30}} \right)$$

30. We have, $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $a\hat{i} + 6\hat{j} - 8\hat{k}$

Two vectors are collinear if and only if,

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} = \lambda \Rightarrow \frac{2}{a} = \frac{-3}{6} = \frac{4}{-8} = \frac{-1}{2} = \lambda$$

$$\Rightarrow \frac{2}{a} = \frac{-1}{2} \Rightarrow a = -4$$

31. $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$

$$|\vec{a}| = \sqrt{(2)^2 + (1)^2 + (2)^2} = \sqrt{4+1+4} = \sqrt{9} = 3.$$

$$\text{Required unit vector is } \hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{2\hat{i} + \hat{j} + 2\hat{k}}{3} \\ = \frac{2}{3}\hat{i} + \frac{1}{3}\hat{j} + \frac{2}{3}\hat{k}$$

32. Refer to answer 31.


33. Refer to answer 31.

34. Position vector which divides the line segment joining points with position vectors $3\vec{a} + \vec{b}$ and $\vec{a} - 3\vec{b}$ in the ratio 2 : 1 externally is given by

$$\frac{2(\vec{a} - 3\vec{b}) - 1(3\vec{a} + \vec{b})}{2-1} = \frac{2\vec{a} - 6\vec{b} - 3\vec{a} - \vec{b}}{1} \\ = -\vec{a} - 7\vec{b}$$

35. Take A to be as origin $(0, 0, 0)$.

\therefore Coordinates of B are $(0, 1, 1)$ and coordinates of C are $(3, -1, 4)$.



Let D be the mid point of BC and AD is a median of $\triangle ABC$.

$$\therefore \text{Coordinates of D are } \left(\frac{3}{2}, 0, \frac{5}{2} \right)$$

$$\text{So, length of AD} = \sqrt{\left(\frac{3}{2} - 0 \right)^2 + (0)^2 + \left(\frac{5}{2} - 0 \right)^2}$$

$$= \sqrt{\frac{9}{4} + \frac{25}{4}} = \frac{\sqrt{34}}{2} \text{ units}$$

36. $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$

$$\therefore \vec{a} + \vec{b} = (2\hat{i} + 3\hat{j} - \hat{k}) + (\hat{i} - 2\hat{j} + \hat{k}) = 3\hat{i} + \hat{j}$$

$$|\vec{a} + \vec{b}| = \sqrt{3^2 + 1^2} = \sqrt{10}$$

∴ A vector of magnitude 5 in the direction of

$$\vec{a} + \vec{b} \text{ is } \frac{5(\vec{a} + \vec{b})}{|\vec{a} + \vec{b}|} = \frac{5(3\hat{i} + \hat{j})}{\sqrt{10}}$$

37. (c) : Here, $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{b} = 2\hat{i} + \lambda\hat{k}$

Since, projection of \vec{a} on $\vec{b} = 0$

$$\Rightarrow \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = 0 \Rightarrow \frac{(\hat{i} - 2\hat{j} + 3\hat{k}) \cdot (2\hat{i} + \lambda\hat{k})}{\sqrt{2^2 + \lambda^2}} = 0$$

$$\Rightarrow \frac{2 + 3\lambda}{\sqrt{4 + \lambda^2}} = 0 \Rightarrow 2 + 3\lambda = 0 \Rightarrow \lambda = -\frac{2}{3}$$

38. Given, two diagonals \vec{d}_1 and \vec{d}_2 are $2\hat{i}$ and $-3\hat{k}$ respectively.

$$\therefore \vec{d}_1 \times \vec{d}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & 0 \\ 0 & 0 & -3 \end{vmatrix} = \hat{i}(0) - \hat{j}(-6 - 0) + \hat{k}(0) = 6\hat{j}$$

$$\begin{aligned} \text{So, area of the parallelogram} &= \frac{1}{2} |\vec{d}_1 \times \vec{d}_2| \\ &= \frac{1}{2} \times 6 = 3 \text{ sq. units} \end{aligned}$$

39. Let $\vec{a} = 2\hat{i} - \lambda\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} - \hat{k}$

We know, \vec{a} and \vec{b} are orthogonal iff $\vec{a} \cdot \vec{b} = 0$

$$\Rightarrow (2\hat{i} - \lambda\hat{j} + \hat{k}) \cdot (\hat{i} + 2\hat{j} - \hat{k}) = 0$$

$$\Rightarrow 2 - 2\lambda - 1 = 0 \Rightarrow 1 - 2\lambda = 0 \Rightarrow \lambda = \frac{1}{2}$$

40. (c) : Since, $\hat{i}, \hat{j}, \hat{k}$ are mutually perpendicular.

$$\therefore \hat{i} \cdot \hat{k} = 0$$

41. Given, $|\vec{a}| = |\vec{b}|$, $\theta = 60^\circ$ and $\vec{a} \cdot \vec{b} = \frac{9}{2}$

$$\text{Now, } \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

$$\Rightarrow \cos 60^\circ = \frac{9/2}{|\vec{a}|^2} \Rightarrow \frac{1}{2} = \frac{9/2}{|\vec{a}|^2}$$

$$\Rightarrow |\vec{a}|^2 = 9 \Rightarrow |\vec{a}| = 3 \therefore |\vec{a}| = |\vec{b}| = 3$$

42. Given, $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = \hat{j} + \hat{k}$

Unit vectors perpendicular to \vec{a} and \vec{b} are

$$\pm \left(\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} \right).$$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 2 \\ 0 & 1 & 1 \end{vmatrix} = -\hat{i} - 2\hat{j} + 2\hat{k}$$

∴ Unit vectors perpendicular to \vec{a} and \vec{b} are

$$\pm \frac{(-\hat{i} - 2\hat{j} + 2\hat{k})}{\sqrt{(-1)^2 + (-2)^2 + (2)^2}} = \pm \left(-\frac{1}{3}\hat{i} - \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k} \right)$$

So, there are two unit vectors perpendicular to the given vectors.

43. We have $\vec{a}, \vec{b}, \vec{c}$ are unit vectors.

Therefore, $|\vec{a}| = 1, |\vec{b}| = 1$ and $|\vec{c}| = 1$

Also, $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ (given)

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 0$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 1 + 1 + 1 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 3 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow (\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = -\frac{3}{2}$$

44. $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 400$

$$\Rightarrow \{|\vec{a}| |\vec{b}| \sin \theta\}^2 + \{|\vec{a}| |\vec{b}| \cos \theta\}^2 = 400$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 \sin^2 \theta + |\vec{a}|^2 |\vec{b}|^2 \cos^2 \theta = 400$$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 = 400 \Rightarrow 25 \times |\vec{b}|^2 = 400 \quad [\because |\vec{a}| = 5]$$

$$\Rightarrow |\vec{b}|^2 = 16 \Rightarrow |\vec{b}| = 4$$

45. Projection of \vec{a} on $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

$$= \frac{(7\hat{i} + \hat{j} - 4\hat{k}) \cdot (2\hat{i} + 6\hat{j} + 3\hat{k})}{\sqrt{(2)^2 + (6)^2 + (3)^2}} = \frac{14 + 6 - 12}{7} = \frac{8}{7}$$

46. Here \hat{a}, \hat{b} and \hat{c} are mutually perpendicular unit vectors.

$$\Rightarrow |\hat{a}| = |\hat{b}| = |\hat{c}| = 1 \text{ and } \hat{a} \cdot \hat{b} = \hat{b} \cdot \hat{c} = \hat{c} \cdot \hat{a} = 0 \quad \dots(1)$$

$$\therefore |2\hat{a} + \hat{b} + \hat{c}|^2 = (2\hat{a} + \hat{b} + \hat{c}) \cdot (2\hat{a} + \hat{b} + \hat{c})$$

$$= 4\hat{a} \cdot \hat{a} + 2\hat{a} \cdot \hat{b} + 2\hat{a} \cdot \hat{c} + 2\hat{b} \cdot \hat{a} + \hat{b} \cdot \hat{b} + \hat{b} \cdot \hat{c} + 2\hat{c} \cdot \hat{a}$$

$$+ \hat{c} \cdot \hat{b} + \hat{c} \cdot \hat{c}$$

$$= 4|\hat{a}|^2 + |\hat{b}|^2 + |\hat{c}|^2 + 4\hat{a} \cdot \hat{b} + 2\hat{b} \cdot \hat{c} + 4\hat{a} \cdot \hat{c}$$

$$(\because \hat{b} \cdot \hat{a} = \hat{a} \cdot \hat{b}, \hat{c} \cdot \hat{a} = \hat{a} \cdot \hat{c}, \hat{c} \cdot \hat{b} = \hat{b} \cdot \hat{c})$$

$$= 4 \cdot 1^2 + 1^2 + 1^2 \quad [\text{Using (1)}]$$

$$= 6$$

$$\therefore |2\hat{a} + \hat{b} + \hat{c}| = \sqrt{6}.$$

47. Here, $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$

Vector perpendicular to both \vec{a} and \vec{b} is

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix} = -\hat{i} + \hat{j} + 0\hat{k} = -\hat{i} + \hat{j}$$

\therefore Unit vector perpendicular to both \vec{a} and \vec{b}

$$= \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{-\hat{i} + \hat{j}}{\sqrt{(-1)^2 + 1^2}} = \frac{1}{\sqrt{2}}(-\hat{i} + \hat{j}).$$

48. Let $\vec{a} = 2\hat{i} - 3\hat{k}$ and $\vec{b} = 4\hat{j} + 2\hat{k}$

The area of a parallelogram with \vec{a} and \vec{b} as its adjacent sides is given by $|\vec{a} \times \vec{b}|$.

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & -3 \\ 0 & 4 & 2 \end{vmatrix} = 12\hat{i} - 4\hat{j} + 8\hat{k}$$

$$\therefore |\vec{a} \times \vec{b}| = \sqrt{(12)^2 + (-4)^2 + (8)^2} = \sqrt{144 + 16 + 64} = \sqrt{224} = 4\sqrt{14} \text{ sq. units.}$$

49. Let θ be the angle between the unit vectors \vec{a} and \vec{b} .

$$\therefore \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \vec{a} \cdot \vec{b} \quad (\because |\vec{a}| = 1 = |\vec{b}|) \dots (1)$$

$$\text{Now } 1 = |\sqrt{2}\vec{a} - \vec{b}|$$

$$\Rightarrow 1 = |\sqrt{2}\vec{a} - \vec{b}|^2 = (\sqrt{2}\vec{a} - \vec{b}) \cdot (\sqrt{2}\vec{a} - \vec{b})$$

$$= 2|\vec{a}|^2 - \sqrt{2}\vec{a} \cdot \vec{b} - \vec{b} \cdot \sqrt{2}\vec{a} + |\vec{b}|^2 = 2 - 2\sqrt{2}\vec{a} \cdot \vec{b} + 1$$

$$(\because \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a})$$

$$= 3 - 2\sqrt{2}\vec{a} \cdot \vec{b}$$

$$\Rightarrow \vec{a} \cdot \vec{b} = \frac{1}{\sqrt{2}} \Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \quad [\text{By using (1)}]$$

$$\therefore \theta = \pi/4$$

50. Refer to answer 45.

51. Refer to answer 45.

52. Given $|\vec{a}| = 1 = |\vec{b}|$, $|\vec{a} + \vec{b}| = 1$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = 1 \Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 1$$

$$\Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = 1$$

$$\Rightarrow |\vec{a}|^2 + 2\vec{a} \cdot \vec{b} + |\vec{b}|^2 = 1 \Rightarrow 1 + 2\vec{a} \cdot \vec{b} + 1 = 1$$

$$\Rightarrow 2\vec{a} \cdot \vec{b} = -1 \Rightarrow 2|\vec{a}| \cdot |\vec{b}| \cos \theta = -1$$

$$\Rightarrow 2 \cdot 1 \cdot 1 \cos \theta = -1$$

$$\Rightarrow \cos \theta = -\frac{1}{2} \Rightarrow \theta = 120^\circ$$

53. Given, $|\vec{a}| = 3$, $|\vec{b}| = \frac{2}{3}$, $|\vec{a} \times \vec{b}| = 1$

$$\Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 1 \Rightarrow 3 \cdot \frac{2}{3} \sin \theta = 1$$

$$\Rightarrow \sin \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}$$

54. Given: $\vec{a} \perp \vec{b} \Rightarrow \vec{a} \cdot \vec{b} = 0$

Also, $|\vec{a}| = 5$ and $|\vec{a} + \vec{b}| = 13$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = 13^2 \Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 169$$

$$\Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = 169$$

$$\Rightarrow |\vec{a}|^2 + 0 + 0 + |\vec{b}|^2 = 169$$

$$\Rightarrow |\vec{b}|^2 = 169 - |\vec{a}|^2 = 169 - 5^2 = 144$$

$$\Rightarrow |\vec{b}| = 12$$

55. The projection of the vector $(\hat{i} + \hat{j} + \hat{k})$ along

$$\text{the vector } \hat{j} \text{ is } (\hat{i} + \hat{j} + \hat{k}) \cdot \left(\frac{\hat{j}}{\sqrt{0^2 + 1^2 + 0^2}} \right) = 1$$

56. We have,

$$\hat{i} \times (\hat{j} + \hat{k}) + \hat{j} \times (\hat{k} + \hat{i}) + \hat{k} \times (\hat{i} + \hat{j})$$

$$= \hat{i} \times \hat{j} + \hat{i} \times \hat{k} + \hat{j} \times \hat{k} + \hat{j} \times \hat{i} + \hat{k} \times \hat{i} + \hat{k} \times \hat{j}$$

$$= \hat{k} - \hat{j} + \hat{i} - \hat{k} + \hat{j} - \hat{i} = \vec{0}.$$

57. Refer to answer 45.

58. Refer to answer 49.

59. Let θ be the angle between the vector

$\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and y -axis i.e., $\vec{b} = \hat{j}$

$$\therefore \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{(\hat{i} + \hat{j} + \hat{k}) \cdot \hat{j}}{|\hat{i} + \hat{j} + \hat{k}| |\hat{j}|}$$

$$= \frac{1}{\sqrt{1^2 + 1^2 + 1^2} \sqrt{1^2}} = \frac{1}{\sqrt{3}}$$

60. Let angle between the vectors \vec{a} and \vec{b} be θ .

Given: $|\vec{a}| = 8$, $|\vec{b}| = 3$ and $|\vec{a} \times \vec{b}| = 12$

$$\Rightarrow |\vec{a}| \cdot |\vec{b}| \cdot \sin \theta = 12 \Rightarrow 8 \times 3 \sin \theta = 12$$

$$\Rightarrow \sin \theta = \frac{12}{24} = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}.$$

61. Here, $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and vector along x -axis is \hat{i} .

\therefore Angle between \vec{a} and \hat{i} is given by

$$\cos \theta = \frac{\vec{a} \cdot \hat{i}}{\sqrt{1^2 + 1^2 + 1^2} \cdot \sqrt{1^2}} = \frac{(\hat{i} + \hat{j} + \hat{k}) \cdot \hat{i}}{\sqrt{3} \cdot 1} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right)$$

62. Here $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 15$, where \vec{a} is unit vector.

$$\Rightarrow \vec{x} \cdot \vec{x} + \vec{x} \cdot \vec{a} - \vec{a} \cdot \vec{x} - \vec{a} \cdot \vec{a} = 15$$

$$\Rightarrow |\vec{x}|^2 - |\vec{a}|^2 = 15 \quad (\because \vec{x} \cdot \vec{a} = \vec{a} \cdot \vec{x})$$

$$\Rightarrow |\vec{x}|^2 - 1 = 15 \quad (\because |\vec{a}| = 1)$$

$$\Rightarrow |\vec{x}|^2 = 16 = 4^2 \Rightarrow |\vec{x}| = 4$$

63. Here, $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + 3\hat{k}$

For \vec{a} is perpendicular to \vec{b} , $\vec{a} \cdot \vec{b} = 0$

$$\Rightarrow (2\hat{i} + \lambda\hat{j} + \hat{k}) \cdot (\hat{i} - 2\hat{j} + 3\hat{k}) = 0$$

$$\Rightarrow 2 \times 1 + \lambda(-2) + 1 \times 3 = 0$$

$$\Rightarrow 2 - 2\lambda + 3 = 0$$

$$\Rightarrow \lambda = \frac{5}{2}$$

64. Refer to answer 63.

65. Here, $\vec{a} = 2\hat{i} - 2\hat{j} + \hat{k}$

$\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$

$$\Rightarrow \vec{b} + \vec{c} = 3\hat{i} + \hat{j} + 2\hat{k}$$

\therefore Projection of $\vec{b} + \vec{c}$ on \vec{a}

$$= \frac{(\vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a}|} = \frac{(3\hat{i} + \hat{j} + 2\hat{k}) \cdot (2\hat{i} - 2\hat{j} + \hat{k})}{|2\hat{i} - 2\hat{j} + \hat{k}|}$$

$$= \frac{3 \times 2 + 1 \times (-2) + 2 \times 1}{\sqrt{2^2 + (-2)^2 + 1^2}} = \frac{6}{3} = 2$$

66. Here, $\vec{a} = \lambda\hat{i} + \hat{j} + 4\hat{k}$, $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$

Given : Projection of \vec{a} on $\vec{b} = 4$

$$\Rightarrow \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = 4 \Rightarrow \frac{(\lambda\hat{i} + \hat{j} + 4\hat{k}) \cdot (2\hat{i} + 6\hat{j} + 3\hat{k})}{|2\hat{i} + 6\hat{j} + 3\hat{k}|} = 4$$

$$\Rightarrow \frac{2\lambda + 6 + 12}{\sqrt{2^2 + 6^2 + 3^2}} = 4$$

$$\Rightarrow 2\lambda + 18 = 4 \times 7$$

$$\Rightarrow 2\lambda = 28 - 18 = 10 \Rightarrow \lambda = 5.$$

67. $(\hat{i} \times \hat{j}) \cdot \hat{k} + \hat{i} \cdot \hat{j} = \hat{k} \cdot \hat{k} + \hat{i} \cdot \hat{j} = 1 + 0 = 1$

68. $(\hat{k} \times \hat{j}) \cdot \hat{i} + \hat{j} \cdot \hat{k} = -\hat{i} \cdot \hat{i} + \hat{j} \cdot \hat{k} = -1 + 0 = -1$

69. $(\hat{k} \times \hat{i}) \cdot \hat{j} + \hat{i} \cdot \hat{k} = \hat{j} \cdot \hat{j} + 0 = 1 + 0 = 1$

70. Let θ be the angle between \vec{a} and \vec{b} , then

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{\sqrt{6}}{\sqrt{3} \cdot 2} \times \frac{\sqrt{3}}{\sqrt{3}}$$

$$= \frac{\sqrt{18}}{3 \times 2} = \frac{3\sqrt{2}}{3 \times 2} = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$$

$$\therefore \theta = \cos^{-1} \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4}$$

71. Let $\vec{a} = \hat{i} - \hat{j}$ and $\vec{b} = \hat{i} + \hat{j}$

$$|\vec{b}| = \sqrt{1^2 + 1^2} = \sqrt{1+1} = \sqrt{2}$$

Also, $\vec{a} \cdot \vec{b} = (\hat{i} - \hat{j}) \cdot (\hat{i} + \hat{j}) = \hat{i} \cdot \hat{i} + \hat{i} \cdot \hat{j} - \hat{j} \cdot \hat{i} - \hat{j} \cdot \hat{j} = 1 - 1 = 0$

$$\therefore \text{Projection of } \vec{a} \text{ on } \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{0}{\sqrt{2}} = 0$$

72. Here, $|\vec{a}| = \sqrt{3}$, $|\vec{b}| = 2$ and angle between \vec{a} and \vec{b} is 60° .

$$\text{Now } \vec{a} \cdot \vec{b} = |\vec{a}| \cdot |\vec{b}| \cdot \cos 60^\circ = \sqrt{3} \times 2 \times \frac{1}{2} = \sqrt{3}.$$

73. Here, $\vec{a} = 5\hat{i} + 6\hat{j} - 2\hat{k}$ and $\vec{b} = 7\hat{i} + 6\hat{j} + 2\hat{k}$

Vector perpendicular to both \vec{a} and \vec{b} is

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 6 & -2 \\ 7 & 6 & 2 \end{vmatrix}$$

$$= \hat{i}(12 + 12) - \hat{j}(10 + 14) + \hat{k}(30 - 42)$$

$$= 24\hat{i} - 24\hat{j} - 12\hat{k} = 12(2\hat{i} - 2\hat{j} - \hat{k})$$

\therefore Unit vector perpendicular to both \vec{a} and \vec{b}

$$= \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{24\hat{i} - 24\hat{j} - 12\hat{k}}{\sqrt{576 + 576 + 144}} = \frac{12(2\hat{i} - 2\hat{j} - \hat{k})}{\sqrt{1296}}$$

$$= \frac{12(2\hat{i} - 2\hat{j} - \hat{k})}{36} = \frac{1}{3}(2\hat{i} - 2\hat{j} - \hat{k})$$

74. For any two non-zero vectors \vec{a} and \vec{b} , we have

$$|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}| \Rightarrow |\vec{a} + \vec{b}|^2 = |\vec{a} - \vec{b}|^2$$

$$\Rightarrow \vec{a}^2 + \vec{b}^2 + 2\vec{a} \cdot \vec{b} = \vec{a}^2 + \vec{b}^2 - 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 4\vec{a} \cdot \vec{b} = 0 \Rightarrow \vec{a} \cdot \vec{b} = 0$$

So, \vec{a} and \vec{b} are perpendicular vectors.

75. Let $A(2\hat{i} - \hat{j} + \hat{k})$, $B(3\hat{i} + 7\hat{j} + \hat{k})$ and

$C(5\hat{i} + 6\hat{j} + 2\hat{k})$

Then, $\vec{AB} = (3-2)\hat{i} + (7+1)\hat{j} + (1-1)\hat{k} = \hat{i} + 8\hat{j}$

$$\overline{AC} = (5-2)\hat{i} + (6+1)\hat{j} + (2-1)\hat{k} = 3\hat{i} + 7\hat{j} + \hat{k}$$

$$\overline{BC} = (5-3)\hat{i} + (6-7)\hat{j} + (2-1)\hat{k} = 2\hat{i} - \hat{j} + \hat{k}$$

Now, angle between \overline{AC} and \overline{BC} is given by

$$\Rightarrow \cos \theta = \frac{\overline{AC} \cdot \overline{BC}}{|\overline{AC}| |\overline{BC}|} = \frac{6-7+1}{\sqrt{9+49+1}\sqrt{4+1+1}}$$

$$\Rightarrow \cos \theta = 0 \Rightarrow AC \perp BC$$

So, A, B, C are the vertices of right angled triangle.

76. Given, $\hat{a} + \hat{b} = \hat{c}$

$$\Rightarrow (\hat{a} + \hat{b}) \cdot (\hat{a} + \hat{b}) = \hat{c} \cdot \hat{c}$$

$$\Rightarrow \hat{a} \cdot \hat{a} + \hat{a} \cdot \hat{b} + \hat{b} \cdot \hat{b} + \hat{b} \cdot \hat{a} = \hat{c} \cdot \hat{c}$$

$$\Rightarrow 1 + \hat{a} \cdot \hat{b} + 1 + \hat{a} \cdot \hat{b} = 1$$

$$\Rightarrow 2 \hat{a} \cdot \hat{b} = -1 \quad \dots(i)$$

Now $(\hat{a} - \hat{b})^2 = (\hat{a} - \hat{b}) \cdot (\hat{a} - \hat{b})$

$$= \hat{a} \cdot \hat{a} - \hat{a} \cdot \hat{b} - \hat{b} \cdot \hat{a} + \hat{b} \cdot \hat{b} = 1 - \hat{a} \cdot \hat{b} - \hat{a} \cdot \hat{b} + 1$$

$$= 2 - 2\hat{a} \cdot \hat{b} = 2 - (-1) \quad [\text{Using(i)}]$$

$$= 3$$

$$\therefore |\hat{a} - \hat{b}| = \sqrt{3}$$

77. Given, $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$ and $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$

Now, $\vec{a} + \vec{b} = 4\hat{i} + \hat{j} - \hat{k}$

Also, $\vec{a} - \vec{b} = -2\hat{i} + 3\hat{j} - 5\hat{k}$

Now, $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = (4\hat{i} + \hat{j} - \hat{k}) \cdot (-2\hat{i} + 3\hat{j} - 5\hat{k})$

$$= (4)(-2) + (1)(3) + (-1)(-5) = -8 + 3 + 5 = 0$$

Hence, $(\vec{a} + \vec{b})$ and $(\vec{a} - \vec{b})$ are perpendicular to each other.

78. Let $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{b} = 3\hat{i} - 2\hat{j} + \hat{k}$

Now, $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$

$$\Rightarrow (\hat{i} - 2\hat{j} + 3\hat{k}) \cdot (3\hat{i} - 2\hat{j} + \hat{k}) = \sqrt{(1)^2 + (-2)^2 + (3)^2} \times \sqrt{(3)^2 + (-2)^2 + (1)^2} \cos \theta$$

$$\Rightarrow 3 + 4 + 3 = \sqrt{14} \times \sqrt{14} \cos \theta$$

$$\Rightarrow \cos \theta = \frac{10}{14}$$


$$\therefore \sin \theta = \sqrt{1 - \cos^2 \theta} = \sqrt{1 - \frac{100}{196}} = \sqrt{\frac{96}{196}}$$

$$\Rightarrow \sin \theta = \frac{4\sqrt{6}}{14} = \frac{2\sqrt{6}}{7}$$

79. Let $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$ and $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$

Then diagonal \overline{AC} of the parallelogram is

$$\begin{aligned} \vec{p} &= \vec{a} + \vec{b} \\ &= \hat{i} + 2\hat{j} + 3\hat{k} + 2\hat{i} + 4\hat{j} - 5\hat{k} \\ &= 3\hat{i} + 6\hat{j} - 2\hat{k} \end{aligned}$$



Therefore unit vector parallel to it is

$$\frac{\vec{p}}{|\vec{p}|} = \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{9+36+4}} = \frac{1}{7}(3\hat{i} + 6\hat{j} - 2\hat{k})$$

Now, diagonal \overline{BD} of the parallelogram is

$$\vec{p}' = \vec{b} - \vec{a} = 2\hat{i} + 4\hat{j} - 5\hat{k} - \hat{i} - 2\hat{j} - 3\hat{k} = \hat{i} + 2\hat{j} - 8\hat{k}$$

Therefore unit vector parallel to it is

$$\frac{\vec{p}'}{|\vec{p}'|} = \frac{\hat{i} + 2\hat{j} - 8\hat{k}}{\sqrt{1+4+64}} = \frac{1}{\sqrt{69}}(\hat{i} + 2\hat{j} - 8\hat{k})$$

80. Given, ΔABC with vertices

$$A(1, 2, 3) \equiv \hat{i} + 2\hat{j} + 3\hat{k}, B(2, -1, 4) \equiv 2\hat{i} - \hat{j} + 4\hat{k},$$

$$C(4, 5, -1) \equiv 4\hat{i} + 5\hat{j} - \hat{k}$$

Now $\overline{AB} = \overline{OB} - \overline{OA} = (2\hat{i} - \hat{j} + 4\hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k}) = \hat{i} - 3\hat{j} + \hat{k}$.

$$\begin{aligned} \overline{AC} &= \overline{OC} - \overline{OA} = (4\hat{i} + 5\hat{j} - \hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k}) \\ &= 3\hat{i} + 3\hat{j} - 4\hat{k}. \end{aligned}$$

$$\therefore (\overline{AB} \times \overline{AC}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 1 \\ 3 & 3 & -4 \end{vmatrix} = 9\hat{i} + 7\hat{j} + 12\hat{k}$$

Hence, area of ΔABC

$$= \frac{1}{2} |\overline{AB} \times \overline{AC}| = \frac{1}{2} |9\hat{i} + 7\hat{j} + 12\hat{k}|$$

$$= \frac{1}{2} \sqrt{9^2 + 7^2 + 12^2} = \frac{1}{2} \sqrt{81 + 49 + 144}$$

$$= \frac{1}{2} \sqrt{274} \text{ sq. units}$$

81. Given, position vector of A = $\hat{i} + \hat{j} + \hat{k}$

Position vector of B = $2\hat{i} + 5\hat{j}$

Position vector of C = $3\hat{i} + 2\hat{j} - 3\hat{k}$

Position vector of D = $\hat{i} - 6\hat{j} - \hat{k}$

$$\therefore \overline{AB} = (2\hat{i} + 5\hat{j}) - (\hat{i} + \hat{j} + \hat{k}) = \hat{i} + 4\hat{j} - \hat{k} \text{ and}$$

$$\overline{CD} = (\hat{i} - 6\hat{j} - \hat{k}) - (3\hat{i} + 2\hat{j} - 3\hat{k}) = -2\hat{i} - 8\hat{j} + 2\hat{k}$$

Now $|\overline{AB}| = \sqrt{(1)^2 + (4)^2 + (1)^2} = \sqrt{18}$

$$|\overline{CD}| = \sqrt{(-2)^2 + (-8)^2 + (2)^2} = \sqrt{4 + 64 + 4} \\ = \sqrt{72} = 2\sqrt{18}$$

Let θ be the angle between \overline{AB} and \overline{CD} .

$$\therefore \cos\theta = \frac{\overline{AB} \cdot \overline{CD}}{|\overline{AB}| |\overline{CD}|} = \frac{(\hat{i} + 4\hat{j} - \hat{k}) \cdot (-2\hat{i} - 8\hat{j} + 2\hat{k})}{(\sqrt{18})(2\sqrt{18})} \\ = \frac{-2 - 32 - 2}{36} = \frac{-36}{36} = -1$$

$$\Rightarrow \cos\theta = -1 \Rightarrow \theta = \pi$$

Since, angle between \overline{AB} and \overline{CD} is 180° .

$\therefore \overline{AB}$ and \overline{CD} are collinear.

82. Let $\vec{d} = x\hat{i} + y\hat{j} + z\hat{k}$

Now, it is given that, \vec{d} is perpendicular to

$$\vec{b} = \hat{i} - 4\hat{j} + 5\hat{k} \text{ and } \vec{c} = 3\hat{i} + \hat{j} - \hat{k}$$

$$\therefore \vec{d} \cdot \vec{b} = 0 \text{ and } \vec{d} \cdot \vec{c} = 0$$

$$\Rightarrow x - 4y + 5z = 0 \quad \dots(i)$$

$$\text{and } 3x + y - z = 0 \quad \dots(ii)$$

Also, $\vec{d} \cdot \vec{a} = 21$, where $\vec{a} = 4\hat{i} + 5\hat{j} - \hat{k}$
 $\Rightarrow 4x + 5y - z = 21 \quad \dots(iii)$

Eliminating z from (i) and (ii), we get
 $16x + y = 0 \quad \dots(iv)$

Eliminating z from (ii) and (iii), we get
 $x + 4y = 21 \quad \dots(v)$

Solving (iv) and (v), we get

$$x = \frac{-1}{3}, y = \frac{16}{3}$$

Putting the values of x and y in (i), we get $z = \frac{13}{3}$

$$\therefore \vec{d} = \frac{-1}{3}\hat{i} + \frac{16}{3}\hat{j} + \frac{13}{3}\hat{k} \text{ is the required vector.}$$

83. $|\vec{a}| = |\vec{b}| = |\vec{c}|$ (Given) $\dots(i)$

and $\vec{a} \cdot \vec{b} = 0, \vec{b} \cdot \vec{c} = 0, \vec{c} \cdot \vec{a} = 0 \quad \dots(ii)$

Let $(\vec{a} + \vec{b} + \vec{c})$ be inclined to vectors $\vec{a}, \vec{b}, \vec{c}$ by angles α, β and γ respectively. Then

$$\cos\alpha = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} = \frac{\vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{a} + \vec{c} \cdot \vec{a}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} \\ = \frac{|\vec{a}|^2 + 0 + 0}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} \quad \text{[Using (ii)]} \\ = \frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|} \quad \dots(iii)$$

Similarly, $\cos\beta = \frac{|\vec{b}|}{|\vec{a} + \vec{b} + \vec{c}|} \quad \dots(iv)$

and $\cos\gamma = \frac{|\vec{c}|}{|\vec{a} + \vec{b} + \vec{c}|} \quad \dots(v)$

From (i), (iii), (iv) and (v), we get
 $\cos\alpha = \cos\beta = \cos\gamma \Rightarrow \alpha = \beta = \gamma$

Hence, the vector $\vec{a} + \vec{b} + \vec{c}$ is equally inclined to the vector \vec{a}, \vec{b} and \vec{c} .

Also the angle between them is given as

$$\alpha = \cos^{-1}\left(\frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|}\right), \beta = \cos^{-1}\left(\frac{|\vec{b}|}{|\vec{a} + \vec{b} + \vec{c}|}\right),$$

$$\gamma = \cos^{-1}\left(\frac{|\vec{c}|}{|\vec{a} + \vec{b} + \vec{c}|}\right)$$

84. We have, $A(2\hat{i} - \hat{j} + \hat{k}), B(\hat{i} - 3\hat{j} - 5\hat{k})$ and $C(3\hat{i} - 4\hat{j} - 4\hat{k})$

Then, $\overline{AB} = (1-2)\hat{i} + (-3+1)\hat{j} + (-5-1)\hat{k}$
 $= -\hat{i} - 2\hat{j} - 6\hat{k}$

$$\overline{AC} = (3-2)\hat{i} + (-4+1)\hat{j} + (-4-1)\hat{k} = \hat{i} - 3\hat{j} - 5\hat{k}$$

$$\text{and } \overline{BC} = (3-1)\hat{i} + (-4+3)\hat{j} + (-4+5)\hat{k} = 2\hat{i} - \hat{j} + \hat{k}$$

Now angle between \overline{AC} and \overline{BC} is given by

$$\cos\theta = \frac{(\overline{AC})(\overline{BC})}{|\overline{AC}| |\overline{BC}|} = \frac{2 + 3 - 5}{\sqrt{1+9+25} \cdot \sqrt{4+1+1}}$$

$$\Rightarrow \cos\theta = 0 \Rightarrow BC \perp AC$$

So, A, B, C are vertices of right angled triangle.


Now area of $\Delta ABC = \frac{1}{2} |\overline{AC} \times \overline{BC}|$

$$= \frac{1}{2} \left\| \begin{matrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & -5 \\ 2 & -1 & 1 \end{matrix} \right\| = \frac{1}{2} |(-3-5)\hat{i} - (1+10)\hat{j} + (-1+6)\hat{k}|$$

$$= \frac{1}{2} |-8\hat{i} - 11\hat{j} + 5\hat{k}|$$

$$= \frac{1}{2} \sqrt{64 + 121 + 25} = \frac{\sqrt{210}}{2} \text{ sq. units.}$$

85. Let $\vec{a} = 2\hat{i} - 4\hat{j} - 5\hat{k}$ and $\vec{b} = 2\hat{i} + 2\hat{j} + 3\hat{k}$



Then diagonal \overline{AC} of the parallelogram is

$$\vec{p} = \vec{a} + \vec{b}$$

$$= 2\hat{i} - 4\hat{j} - 5\hat{k} + 2\hat{i} + 2\hat{j} + 3\hat{k} = 4\hat{i} - 2\hat{j} - 2\hat{k}$$

Therefore, unit vector parallel to it is

$$\frac{\vec{p}}{|\vec{p}|} = \frac{4\hat{i} - 2\hat{j} - 2\hat{k}}{\sqrt{16+4+4}} = \frac{2\hat{i} - \hat{j} - \hat{k}}{\sqrt{6}}$$

Now, diagonal \overline{BD} of the parallelogram is

$$\vec{p}' = \vec{b} - \vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k} - 2\hat{i} + 4\hat{j} + 5\hat{k} = 6\hat{j} + 8\hat{k}$$

Therefore, unit vector parallel to it is

$$\frac{\vec{p}'}{|\vec{p}'|} = \frac{6\hat{j} + 8\hat{k}}{\sqrt{36+64}} = \frac{6\hat{j} + 8\hat{k}}{10} = \frac{3\hat{j} + 4\hat{k}}{5}$$

$$\text{Now, } \vec{p} \times \vec{p}' = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & -2 & -2 \\ 0 & 6 & 8 \end{vmatrix}$$

$$= \hat{i}(-16+12) - \hat{j}(32-0) + \hat{k}(24-0)$$

$$= -4\hat{i} - 32\hat{j} + 24\hat{k}$$

$$\therefore \text{Area of parallelogram} = \frac{|\vec{p} \times \vec{p}'|}{2}$$

$$= \frac{\sqrt{16+1024+576}}{2} = 2\sqrt{101} \text{ sq. units.}$$

86. Two non zero vectors are parallel if and only if their cross product is zero vector.

So, we have to prove that cross product of $\vec{a} - \vec{d}$ and $\vec{b} - \vec{c}$ is zero vector.

$$(\vec{a} - \vec{d}) \times (\vec{b} - \vec{c}) = (\vec{a} \times \vec{b}) - (\vec{a} \times \vec{c}) - (\vec{d} \times \vec{b}) + (\vec{d} \times \vec{c})$$

Since, it is given that $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$ and $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$

$$\text{And, } \vec{d} \times \vec{b} = -\vec{b} \times \vec{d}, \vec{d} \times \vec{c} = -\vec{c} \times \vec{d}$$

Therefore,

$$(\vec{a} - \vec{d}) \times (\vec{b} - \vec{c}) = (\vec{c} \times \vec{d}) - (\vec{b} \times \vec{d}) + (\vec{b} \times \vec{d}) - (\vec{c} \times \vec{d}) = \vec{0}$$

Hence, $\vec{a} - \vec{d}$ is parallel to $\vec{b} - \vec{c}$, where

$$\vec{a} \neq \vec{d} \text{ and } \vec{b} \neq \vec{c}.$$

87. $(\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j}) + xy$

$$= [(x\hat{i} + y\hat{j} + z\hat{k}) \times \hat{i}] \cdot [(x\hat{i} + y\hat{j} + z\hat{k}) \times \hat{j}] + xy$$

$$= (-y\hat{k} + z\hat{j}) \cdot (x\hat{k} - z\hat{i}) + xy = -xy + xy = 0$$

88. Here, $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j}$, $\vec{c} = 3\hat{i} - 4\hat{j} - 5\hat{k}$

$$\therefore \vec{a} - \vec{b} = (\hat{i} + 2\hat{j} + \hat{k}) - (2\hat{i} + \hat{j}) = -\hat{i} + \hat{j} + \hat{k}$$

$$\vec{c} - \vec{b} = (3\hat{i} - 4\hat{j} - 5\hat{k}) - (2\hat{i} + \hat{j}) = \hat{i} - 5\hat{j} - 5\hat{k}$$

Vector perpendicular to both $\vec{a} - \vec{b}$ and $\vec{c} - \vec{b}$ is

$$(\vec{a} - \vec{b}) \times (\vec{c} - \vec{b}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & 1 \\ 1 & -5 & -5 \end{vmatrix}$$

$$= (-5+5)\hat{i} - (5-1)\hat{j} + (5-1)\hat{k} = -4\hat{j} + 4\hat{k}$$

\therefore Unit vector perpendicular to both $\vec{a} - \vec{b}$ and $\vec{c} - \vec{b}$

$$= \frac{-4\hat{j} + 4\hat{k}}{|-4\hat{j} + 4\hat{k}|} = \frac{-4\hat{j} + 4\hat{k}}{\sqrt{(-4)^2 + 4^2}} = \frac{-4\hat{j} + 4\hat{k}}{4\sqrt{2}} = \frac{1}{\sqrt{2}}(-\hat{j} + \hat{k}).$$

89. Given $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ and $|\vec{a}| = 3$, $|\vec{b}| = 5$, $|\vec{c}| = 7$

We have $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\Rightarrow \vec{a} + \vec{b} = -\vec{c} \Rightarrow |\vec{a} + \vec{b}|^2 = |-\vec{c}|^2$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2(\vec{a} \cdot \vec{b}) = |\vec{c}|^2$$

$$\Rightarrow 9 + 25 + 2|\vec{a}||\vec{b}|\cos\theta = 49$$

$$\Rightarrow 2 \times 3 \times 5 \times \cos\theta = 49 - 34 = 15$$

$$\Rightarrow \cos\theta = \frac{15}{30} = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3} = 60^\circ$$

90. Here, $\vec{a} = \hat{i} + \hat{j} + \hat{k}$; $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$ and

$$\vec{c} = \lambda\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\Rightarrow \vec{b} + \vec{c} = (2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}$$

The unit vector along $\vec{b} + \vec{c}$ is $\vec{p} = \frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|}$

$$= \frac{(2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{(2 + \lambda)^2 + 6^2 + (-2)^2}} = \frac{(2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{\lambda^2 + 4\lambda + 44}}$$

Also, $\vec{a} \cdot \vec{p} = 1$ (Given)

$$\Rightarrow \frac{(2 + \lambda) + 6 - 2}{\sqrt{\lambda^2 + 4\lambda + 44}} = 1$$

$$\Rightarrow \sqrt{\lambda^2 + 4\lambda + 44} = \lambda + 6$$

$$\Rightarrow \lambda^2 + 4\lambda + 44 = \lambda^2 + 12\lambda + 36$$

$$\Rightarrow 8\lambda = 8 \Rightarrow \lambda = 1$$

\therefore The required unit vector

$$\vec{p} = \frac{(2+1)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{1+4+44}} = \frac{1}{7}(3\hat{i} + 6\hat{j} - 2\hat{k}).$$

91. We have $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$

$$\text{Let } \vec{r} = \vec{a} + \vec{b} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\text{and } \vec{p} = \vec{a} - \vec{b} = -\hat{j} - 2\hat{k}$$

A unit vector perpendicular to both \vec{r} and \vec{p} is

$$\text{given as } \pm \frac{\vec{r} \times \vec{p}}{|\vec{r} \times \vec{p}|}.$$

$$\text{Now, } \vec{r} \times \vec{p} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 0 & -1 & -2 \end{vmatrix} = -2\hat{i} + 4\hat{j} - 2\hat{k}$$

So, the required unit vector is

$$= \pm \frac{(-2\hat{i} + 4\hat{j} - 2\hat{k})}{\sqrt{(-2)^2 + 4^2 + (-2)^2}} = \mp \frac{(\hat{i} - 2\hat{j} + \hat{k})}{\sqrt{6}}$$

92. Here, $\vec{a} = 2\hat{i} - 3\hat{j} + \hat{k}$; $\vec{b} = -\hat{i} + \hat{k}$; $\vec{c} = 2\hat{j} - \hat{k}$

$$\therefore \vec{a} + \vec{b} = (2\hat{i} - 3\hat{j} + \hat{k}) + (-\hat{i} + \hat{k}) = \hat{i} - 3\hat{j} + 2\hat{k},$$

$$\vec{b} + \vec{c} = (-\hat{i} + \hat{k}) + (2\hat{j} - \hat{k}) = -\hat{i} + 2\hat{j}$$

$$\therefore (\vec{a} + \vec{b}) \times (\vec{b} + \vec{c}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ -1 & 2 & 0 \end{vmatrix} = -4\hat{i} - 2\hat{j} - \hat{k}$$

\therefore Area of a parallelogram whose diagonals are $\vec{a} + \vec{b}$ and $\vec{b} + \vec{c}$

$$= \frac{1}{2} |(\vec{a} + \vec{b}) \times (\vec{b} + \vec{c})| = \frac{1}{2} |-4\hat{i} - 2\hat{j} - \hat{k}|$$

$$= \frac{1}{2} \sqrt{(-4)^2 + (-2)^2 + (-1)^2} = \frac{\sqrt{21}}{2} \text{ sq.units.}$$

93. Refer to answer 82.

94. Here $|\vec{a} + \vec{b}| = |\vec{a}|$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = |\vec{a}|^2 \Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = \vec{a} \cdot \vec{a}$$

$$\Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = \vec{a} \cdot \vec{a}$$

$$\Rightarrow 2\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{b} = 0 \quad [\because \vec{b} \cdot \vec{a} = \vec{a} \cdot \vec{b}]$$

$$\Rightarrow (2\vec{a} + \vec{b}) \cdot \vec{b} = 0 \Rightarrow (2\vec{a} + \vec{b}) \perp \vec{b}$$

95. Given $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = \hat{j} - \hat{k}$

Let $\vec{c} = x\hat{i} + y\hat{j} + z\hat{k}$

Now we have, $\vec{a} \times \vec{c} = \vec{b}$

$$\Rightarrow (\hat{i} + \hat{j} + \hat{k}) \times (x\hat{i} + y\hat{j} + z\hat{k}) = \hat{j} - \hat{k}$$

$$\Rightarrow \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ x & y & z \end{vmatrix} = \hat{j} - \hat{k}$$

$$\Rightarrow \hat{i}(z - y) - \hat{j}(z - x) + \hat{k}(y - x) = \hat{j} - \hat{k}$$

$$\Rightarrow z - y = 0, x - z = 1 \text{ and } y - x = -1$$

$$\Rightarrow y = z, x - z = 1, x - y = 1 \quad \dots(i)$$

Also, we have $\vec{a} \cdot \vec{c} = 3$

$$\Rightarrow (\hat{i} + \hat{j} + \hat{k}) \cdot (x\hat{i} + y\hat{j} + z\hat{k}) = 3$$

$$\Rightarrow x + y + z = 3$$

$$\Rightarrow x + x - 1 + x - 1 = 3 \quad [\text{Using (i)}]$$

$$\Rightarrow 3x - 2 = 3 \Rightarrow x = \frac{5}{3}, y = \frac{2}{3}, z = \frac{2}{3}$$

$$\text{Hence, } \vec{c} = \frac{5}{3}\hat{i} + \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k}$$

96. Here $\vec{a} = \hat{i} - \hat{j} + 7\hat{k}$; $\vec{b} = 5\hat{i} - \hat{j} + \lambda\hat{k}$

$$\therefore \vec{a} + \vec{b} = 6\hat{i} - 2\hat{j} + (7 + \lambda)\hat{k}$$

$$\vec{a} - \vec{b} = -4\hat{i} + (7 - \lambda)\hat{k}$$

For $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ to be perpendicular,

$$(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 0$$

$$\Rightarrow [6\hat{i} - 2\hat{j} + (7 + \lambda)\hat{k}] \cdot [-4\hat{i} + (7 - \lambda)\hat{k}] = 0$$

$$\Rightarrow 6 \times (-4) + (7 + \lambda) \times (7 - \lambda) = 0$$

$$\Rightarrow -24 + 49 - \lambda^2 = 0 \Rightarrow \lambda^2 = 25 \Rightarrow \lambda = \pm 5$$

97. Refer to answer 83.

98. Let the required vector be $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$.

Also let,

$$\vec{a} = \hat{i} - \hat{j} + \hat{k}, \vec{b} = 2\hat{i} + \hat{j} - 3\hat{k} \text{ and } \vec{c} = \hat{i} + \hat{j} + \hat{k}$$

$$\vec{r} \cdot \vec{a} = 4, \vec{r} \cdot \vec{b} = 0, \vec{r} \cdot \vec{c} = 2 \quad (\text{Given})$$

$$\Rightarrow x - y + z = 4 \quad \dots(i)$$

$$2x + y - 3z = 0 \quad \dots(ii)$$

$$x + y + z = 2 \quad \dots(iii)$$

$$\text{Now (iii) - (i)} \Rightarrow 2y = -2 \Rightarrow y = -1$$

From (ii) and (iii)

$$2x - 3z - 1 = 0, x + z - 3 = 0 \Rightarrow x = 2, z = 1$$

\therefore The required vector is $\vec{r} = 2\hat{i} - \hat{j} + \hat{k}$.

99. Here, $\vec{a} = 2\lambda^2\hat{i} + 4\lambda\hat{j} + \hat{k}$ and

$$\vec{b} = 7\hat{i} - 2\hat{j} + \lambda\hat{k}$$

If θ is the angle between the vectors \vec{a} and \vec{b} ,

$$\text{then } \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

For θ to be obtuse, $\cos\theta < 0 \Rightarrow \vec{a} \cdot \vec{b} < 0$

$$\Rightarrow (2\lambda^2\hat{i} + 4\lambda\hat{j} + \hat{k}) \cdot (7\hat{i} - 2\hat{j} + \lambda\hat{k}) < 0$$

$$\Rightarrow 2\lambda^2 \cdot 7 + 4\lambda \cdot (-2) + 1 \cdot \lambda < 0$$

$$\Rightarrow 14\lambda^2 - 7\lambda < 0 \Rightarrow \lambda(2\lambda - 1) < 0$$

$$\Rightarrow \text{Either } \lambda < 0, 2\lambda - 1 > 0 \text{ or } \lambda > 0, 2\lambda - 1 < 0$$

$$\Rightarrow \text{Either } \lambda < 0, \lambda > \frac{1}{2} \text{ or } \lambda > 0, \lambda < \frac{1}{2}$$

First alternative is impossible.

$$\therefore \lambda > 0, \lambda < \frac{1}{2} \text{ i.e., } 0 < \lambda < \frac{1}{2} \text{ i.e., } \lambda \in \left] 0, \frac{1}{2} \right[$$

100. Given, $|\vec{a}| = 3, |\vec{b}| = 4, |\vec{c}| = 5 \quad \dots(i)$

$$\text{and } \vec{a} \cdot (\vec{b} + \vec{c}) = 0, \vec{b} \cdot (\vec{c} + \vec{a}) = 0, \vec{c} \cdot (\vec{a} + \vec{b}) = 0$$

$$\begin{aligned} \therefore \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c} + \vec{b} \cdot \vec{a} + \vec{c} \cdot \vec{a} + \vec{c} \cdot \vec{b} \\ = 0 + 0 + 0 = 0 \\ \Rightarrow 2(\vec{a} \cdot \vec{b}) + 2(\vec{b} \cdot \vec{c}) + 2(\vec{c} \cdot \vec{a}) = 0 \quad \dots(\text{ii}) \\ \text{Now } |\vec{a} + \vec{b} + \vec{c}|^2 = (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) \\ = (\vec{a} + \vec{b} + \vec{c}) \cdot \vec{a} + (\vec{a} + \vec{b} + \vec{c}) \cdot \vec{b} + (\vec{a} + \vec{b} + \vec{c}) \cdot \vec{c} \\ = \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{a} + \vec{c} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{b} + \vec{c} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c} \\ + \vec{c} \cdot \vec{c} \\ = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b}) + 2(\vec{b} \cdot \vec{c}) + 2(\vec{c} \cdot \vec{a}) \\ = 3^2 + 4^2 + 5^2 + 0 \quad [\text{Using (i) and (ii)}] \\ = 50 \end{aligned}$$

$$\therefore |\vec{a} + \vec{b} + \vec{c}| = 5\sqrt{2}.$$

101. Here $\vec{a} = 3\hat{i} - \hat{j}$, $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$
We have to express: $\vec{b} = \vec{b}_1 + \vec{b}_2$, where
 $\vec{b}_1 \parallel \vec{a}$ and $\vec{b}_2 \perp \vec{a}$
Let $\vec{b}_1 = \lambda \vec{a} = \lambda(3\hat{i} - \hat{j})$ and $\vec{b}_2 = x\hat{i} + y\hat{j} + z\hat{k}$
Now $\vec{b}_2 \perp \vec{a} \Rightarrow \vec{b}_2 \cdot \vec{a} = 0$
 $\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} - \hat{j}) = 0$
 $\Rightarrow 3x - y = 0 \quad \dots(\text{i})$

Now, $\vec{b} = \vec{b}_1 + \vec{b}_2$
 $\Rightarrow 2\hat{i} + \hat{j} - 3\hat{k} = \lambda(3\hat{i} - \hat{j}) + (x\hat{i} + y\hat{j} + z\hat{k})$
On comparing, we get
$$\left. \begin{aligned} 2 &= 3\lambda + x \\ 1 &= -\lambda + y \end{aligned} \right\} \Rightarrow x + 3y = 5 \quad \dots(\text{ii})$$

and $-3 = z \Rightarrow z = -3$
Solving (i) and (ii), we get $x = \frac{1}{2}$, $y = \frac{3}{2}$

$$\therefore 1 = -\lambda + y = -\lambda + \frac{3}{2} \Rightarrow \lambda = \frac{1}{2}$$

Hence, $\vec{b}_1 = \lambda(3\hat{i} - \hat{j}) = \frac{3}{2}\hat{i} - \frac{1}{2}\hat{j}$

and $\vec{b}_2 = \frac{1}{2}\hat{i} + \frac{3}{2}\hat{j} - 3\hat{k}$

102. We have,
 $|\vec{a}| = 5$, $|\vec{b}| = 12$, $|\vec{c}| = 13$ and $\vec{a} + \vec{b} + \vec{c} = \vec{0}$
 $\Rightarrow (\vec{a} + \vec{b} + \vec{c})^2 = |\vec{0}|^2$ (Squaring on both sides)
 $\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = 0$
 $\Rightarrow 25 + 144 + 169 + 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = 0$
 $\Rightarrow 2[\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}] = -338$
 $\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = \frac{-338}{2} = -169$

103. Refer to answer 82.

104. Refer to answer 76.

105. Refer to answer 91.

106. We have $|\vec{a}| = 2$, $|\vec{b}| = 1$ and $\vec{a} \cdot \vec{b} = 1$
Now, $(3\vec{a} - 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$
 $= 6|\vec{a}|^2 + 21\vec{a} \cdot \vec{b} - 10\vec{a} \cdot \vec{b} - 35|\vec{b}|^2$
 $= 6|\vec{a}|^2 + 11\vec{a} \cdot \vec{b} - 35|\vec{b}|^2$
 $= 6(2)^2 + 11(1) - 35(1)^2 = 24 + 11 - 35 = 0$

107. Refer to answer 80.

108. Refer to answer 97.
Also the angle between them is given as

$$\alpha = \cos^{-1} \left(\frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|} \right),$$

$$\beta = \cos^{-1} \left(\frac{|\vec{b}|}{|\vec{a} + \vec{b} + \vec{c}|} \right),$$

$$\gamma = \cos^{-1} \left(\frac{|\vec{c}|}{|\vec{a} + \vec{b} + \vec{c}|} \right)$$

109. Refer to answer 100.

110. Since the vectors are coplanar.

$$\therefore \begin{vmatrix} 1 & 3 & 1 \\ 2 & -1 & -1 \\ 0 & \lambda & 3 \end{vmatrix} = 0$$

$$\Rightarrow 1(-3 + \lambda) - 3(6 - 0) + 1(2\lambda - 0) = 0$$

$$\Rightarrow -3 + \lambda - 18 + 2\lambda = 0$$

$$\Rightarrow 3\lambda - 21 = 0 \Rightarrow \lambda = 7$$

111. Here $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$,
 $\vec{c} = 3\hat{i} + \hat{j} + 2\hat{k}$

Now, $\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 2 & 1 \\ 3 & 1 & 2 \end{vmatrix} = 3\hat{i} + 5\hat{j} - 7\hat{k}$

$$\begin{aligned} \therefore \vec{a} \cdot (\vec{b} \times \vec{c}) &= (2\hat{i} + \hat{j} + 3\hat{k}) \cdot (3\hat{i} + 5\hat{j} - 7\hat{k}) \\ &= 2 \times 3 + 1 \times 5 + 3 \times (-7) \\ &= 6 + 5 - 21 = -10 \end{aligned}$$

112. Given, $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} + 4\hat{j} - 5\hat{k}$ and
 $\vec{c} = 2\hat{i} - \hat{j} + 3\hat{k}$

$$\begin{aligned} \therefore 2\vec{a} &= 2\hat{i} - 2\hat{j} + 4\hat{k} \\ -\vec{b} &= -3\hat{i} - 4\hat{j} + 5\hat{k} \\ 3\vec{c} &= 6\hat{i} - 3\hat{j} + 9\hat{k} \end{aligned}$$

$$\begin{aligned} \text{Now, } 2\vec{a} \cdot (-\vec{b} \times 3\vec{c}) &= \begin{vmatrix} 2 & -2 & 4 \\ -3 & -4 & 5 \\ 6 & -3 & 9 \end{vmatrix} \\ &= 2(-36 + 15) + 2(-27 - 30) + 4(9 + 24) \\ &= 2(-21) - 2(57) + 4(33) \\ &= -42 - 114 + 132 = -24 \end{aligned}$$

\therefore Volume of parallelepiped

$$|2\vec{a} \cdot (-\vec{b} \times 3\vec{c})| = |-24| = 24 \text{ cubic units}$$

113. Given, $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$ and

$$\vec{c} = -3\hat{i} + \hat{j} + 2\hat{k}$$

$$\begin{aligned} \text{Now, } \vec{b} \times \vec{c} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 1 \\ -3 & 1 & 2 \end{vmatrix} \\ &= \hat{i}(-4 - 1) - \hat{j}(2 + 3) + \hat{k}(1 - 6) \\ &= -5\hat{i} - 5\hat{j} - 5\hat{k} \end{aligned}$$

$$\begin{aligned} \therefore \vec{a} \cdot (\vec{b} \times \vec{c}) &= (2\hat{i} + 3\hat{j} + \hat{k}) \cdot (-5\hat{i} - 5\hat{j} - 5\hat{k}) \\ &= -10 - 15 - 5 = -30 \end{aligned}$$

114. Given points are $A(x, -1, -1)$, $B(4, 5, 1)$, $C(3, 9, 4)$ and $D(-4, 4, 4)$.

$$\begin{aligned} \overline{AB} &= (4-x)\hat{i} + (5+1)\hat{j} + (1+1)\hat{k} \\ &= (4-x)\hat{i} + 6\hat{j} + 2\hat{k} \end{aligned}$$

$$\begin{aligned} \overline{AC} &= (3-x)\hat{i} + (9+1)\hat{j} + (4+1)\hat{k} \\ &= (3-x)\hat{i} + 10\hat{j} + 5\hat{k} \end{aligned}$$

$$\begin{aligned} \overline{AD} &= (-4-x)\hat{i} + (4+1)\hat{j} + (4+1)\hat{k} \\ &= -(4+x)\hat{i} + 5\hat{j} + 5\hat{k} \end{aligned}$$

The given points will be coplanar iff

$$[\overline{AB} \ \overline{AC} \ \overline{AD}] = 0$$

$$\text{Now, } [\overline{AB} \ \overline{AC} \ \overline{AD}] = 0 \Rightarrow \begin{vmatrix} 4-x & 6 & 2 \\ 3-x & 10 & 5 \\ -(4+x) & 5 & 5 \end{vmatrix} = 0$$

$$\Rightarrow (4-x)(50-25) - 6(15-5x+20+5x) + 2(15-5x+40+10x) = 0$$

$$\Rightarrow (4-x)(25) - 6(35) + 2(55+5x) = 0$$

$$\Rightarrow 100 - 25x - 210 + 110 + 10x = 0$$

$$\Rightarrow -15x = 0$$

$$\Rightarrow x = 0$$

115. We have, $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i}$, $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$

$$(a) \quad c_1 = 1, c_2 = 2 \quad \therefore \vec{c} = \hat{i} + 2\hat{j} + c_3\hat{k}$$

Given that \vec{a} , \vec{b} and \vec{c} are coplanar

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 2 & c_3 \end{vmatrix} = 0$$

$$\Rightarrow -1(c_3) + 1(2) = 0 \Rightarrow c_3 = 2$$

$$(b) \quad c_2 = -1, c_3 = 1 \quad \therefore \vec{c} = c_1\hat{i} - \hat{j} + \hat{k}$$

Let \vec{a} , \vec{b} and \vec{c} are coplanar.

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ c_1 & -1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -1(1) + 1(-1) = 0 \Rightarrow -2 = 0, \text{ which is false.}$$

So, no value of c_1 can make \vec{a} , \vec{b} and \vec{c} coplanar.

116. Let A, B, C, D be the given points. The given points will be coplanar iff any one of the following triads of vectors are coplanar.

$$\overline{AB}, \overline{AC}, \overline{AD}; \overline{BC}, \overline{BA}, \overline{BD} \text{ etc.}$$

If $\overline{AB}, \overline{AC}, \overline{AD}$ are coplanar, then their scalar triple product $[\overline{AB} \ \overline{AC} \ \overline{AD}] = 0$ where, $A(3\hat{i} + 6\hat{j} + 9\hat{k})$,

$$B(\hat{i} + 2\hat{j} + 3\hat{k}), C(2\hat{i} + 3\hat{j} + \hat{k}) \text{ and } D(4\hat{i} + 6\hat{j} + \lambda\hat{k}).$$

$$\begin{aligned} \text{Now, } \overline{AB} &= (\hat{i} + 2\hat{j} + 3\hat{k}) - (3\hat{i} + 6\hat{j} + 9\hat{k}) \\ &= -2\hat{i} - 4\hat{j} - 6\hat{k} \end{aligned}$$

$$\overline{AC} = (2\hat{i} + 3\hat{j} + \hat{k}) - (3\hat{i} + 6\hat{j} + 9\hat{k}) = -\hat{i} - 3\hat{j} - 8\hat{k}$$

$$\overline{AD} = (4\hat{i} + 6\hat{j} + \lambda\hat{k}) - (3\hat{i} + 6\hat{j} + 9\hat{k}) = \hat{i} + (\lambda - 9)\hat{k}$$

$$\therefore [\overline{AB} \ \overline{AC} \ \overline{AD}] = \begin{vmatrix} -2 & -4 & -6 \\ -1 & -3 & -8 \\ 1 & 0 & \lambda - 9 \end{vmatrix} = 0$$

$$\Rightarrow -2(-3\lambda + 27) + 4(-\lambda + 9 + 8) - 6(0 + 3) = 0$$

$$\Rightarrow 2(3\lambda - 27) - 4(\lambda - 17) - 6(3) = 0$$

$$\Rightarrow 6\lambda - 54 - 4\lambda + 68 - 18 = 0$$

$$\Rightarrow 2\lambda - 4 = 0 \Rightarrow \lambda = 2$$

117. Since, $\vec{a} + \vec{b}$, $\vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$ are coplanar.

$$\therefore (\vec{a} + \vec{b}) \cdot [(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})] = 0$$

$$\Rightarrow (\vec{a} + \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}] = 0$$

$$\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{a}) = 0 \quad [\because \vec{c} \times \vec{c} = 0]$$

$$\Rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) + \vec{b} \cdot (\vec{b} \times \vec{c})$$

$$+ \vec{b} \cdot (\vec{b} \times \vec{a}) + \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$$

$$\Rightarrow 2[\vec{a} \cdot (\vec{b} \times \vec{c})] = 0 \Rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) = 0$$

$\Rightarrow \vec{a}, \vec{b}$ and \vec{c} are coplanar.

118. Here position vectors of A, B, C and D are $4\hat{i} + 5\hat{j} + \hat{k}$, $-\hat{j} - \hat{k}$, $3\hat{i} + \lambda\hat{j} + 4\hat{k}$ and $-4\hat{i} + 4\hat{j} + 4\hat{k}$ respectively.

$$\Rightarrow \overline{AB} = -\hat{j} - \hat{k} - (4\hat{i} + 5\hat{j} + \hat{k}) = -4\hat{i} - 6\hat{j} - 2\hat{k}$$

$$\overline{AC} = (3\hat{i} + \lambda\hat{j} + 4\hat{k}) - (4\hat{i} + 5\hat{j} + \hat{k}) = -\hat{i} + (\lambda - 5)\hat{j} + 3\hat{k}$$

$$\overline{AD} = (-4\hat{i} + 4\hat{j} + 4\hat{k}) - (4\hat{i} + 5\hat{j} + \hat{k}) = -8\hat{i} - \hat{j} + 3\hat{k}$$

\therefore For points A, B, C, D to be coplanar

\Leftrightarrow Vectors $\overline{AB}, \overline{AC}, \overline{AD}$ will be coplanar

$$\Leftrightarrow [\overline{AB} \overline{AC} \overline{AD}] = 0$$

$$\Rightarrow \begin{vmatrix} -4 & -6 & -2 \\ -1 & \lambda - 5 & 3 \\ -8 & -1 & 3 \end{vmatrix} = 0$$

$$\Rightarrow -4(3\lambda - 15 + 3) + 6(-3 + 24) - 2(1 + 8\lambda - 40) = 0$$

$$\Rightarrow -12\lambda + 48 + 126 + 78 - 16\lambda = 0$$

$$\Rightarrow 28\lambda = 252 \Rightarrow \lambda = 9.$$

119. We know that $[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c})$

$$\therefore \text{L.H.S.} = [\vec{a}, \vec{b} + \vec{c}, \vec{d}] = \vec{a} \cdot [(\vec{b} + \vec{c}) \times \vec{d}]$$

$$= \vec{a} \cdot [\vec{b} \times \vec{d} + \vec{c} \times \vec{d}] = \vec{a} \cdot (\vec{b} \times \vec{d}) + \vec{a} \cdot (\vec{c} \times \vec{d})$$

$$= [\vec{a}, \vec{b}, \vec{d}] + [\vec{a}, \vec{c}, \vec{d}] = \text{R.H.S.}$$

120. We have, $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}]$

$$= (\vec{a} + \vec{b}) \cdot \{(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})\}$$

$$= (\vec{a} + \vec{b}) \cdot \{(\vec{b} \times \vec{c}) + (\vec{b} \times \vec{a}) + (\vec{c} \times \vec{c}) + (\vec{c} \times \vec{a})\}$$

$$= (\vec{a} + \vec{b}) \cdot \{(\vec{b} \times \vec{c}) + (\vec{b} \times \vec{a}) + (\vec{c} \times \vec{a})\} \quad [\because \vec{c} \times \vec{c} = 0]$$

$$= \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) +$$

$$\vec{b} \cdot (\vec{b} \times \vec{c}) + \vec{b} \cdot (\vec{b} \times \vec{a}) + \vec{b} \cdot (\vec{c} \times \vec{a})$$

$$= [\vec{a} \vec{b} \vec{c}] + [\vec{b} \vec{c} \vec{a}]$$

[\because Scalar triple product with two equal vectors is 0]

$$= [\vec{a} \vec{b} \vec{c}] + [\vec{a} \vec{b} \vec{c}] \quad (\because [\vec{b} \vec{c} \vec{a}] = [\vec{a} \vec{b} \vec{c}])$$

$$= 2[\vec{a} \vec{b} \vec{c}]$$

121. Refer to answer 118.

122. If the vectors $\vec{a} + \vec{b}, \vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$

are coplanar, then $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = 0$

$$\Leftrightarrow (\vec{a} + \vec{b}) \cdot [(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})] = 0$$

$$\Leftrightarrow (\vec{a} + \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}] = 0$$

$$\Leftrightarrow (\vec{a} + \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{0} + \vec{c} \times \vec{a}] = 0$$

$$\Leftrightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a})$$

$$+ \vec{b} \cdot (\vec{b} \times \vec{c}) + \vec{b} \cdot (\vec{b} \times \vec{a}) + \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$$

$$\Leftrightarrow [\vec{a} \vec{b} \vec{c}] + [\vec{a} \vec{b} \vec{a}] + [\vec{a} \vec{c} \vec{a}] +$$

$$[\vec{b} \vec{b} \vec{c}] + [\vec{b} \vec{b} \vec{a}] + [\vec{b} \vec{c} \vec{a}] = 0$$

$$\Leftrightarrow [\vec{a} \vec{b} \vec{c}] + 0 + 0 + 0 + 0 + [\vec{a} \vec{b} \vec{c}] = 0$$

$$\Leftrightarrow 2[\vec{a} \vec{b} \vec{c}] = 0 \Leftrightarrow [\vec{a} \vec{b} \vec{c}] = 0$$

\therefore The vectors $\vec{a}, \vec{b}, \vec{c}$ are coplanar.

Hence the vectors $\vec{a}, \vec{b}, \vec{c}$ are coplanar if and only

if $\vec{a} + \vec{b}, \vec{b} + \vec{c}$ and $\vec{c} + \vec{a}$ are coplanar.

123. Refer to answer 122.